References

1. Sacco RL, Bello JA, Traub R, Brust JC. Selective proprioceptive loss from a thalamic lacunar stroke. *Stroke* 1987;18:1160–1163.
2. Johansson RS, Vallbo AB. Tactile sensory coding in the glabrous skin of the human hand. *Trends Neurosci* 1983;6:27–32.
3. Dykes RW, Sur M, Merzenich MM, Kaas JH, Nelson RJ. Regional segregation of neurons responding to quickly adapting, SA, deep and pacinian receptors within thalamic ventroposterior lateral and ventroposterior inferior nuclei in the squirrel monkey (*Saimiri sciureus*). *Neuroscience* 1981;6:1687–1692.
4. Jones EG, Friedman DP, Hendry SHC. Thalamic basis of place- and modality-specific columns in monkey somatosensory cortex: a correlative anatomical and physiological study. *J Neurophysiol* 1982;48:545–568.
5. Lenz FA, Dostrovsky JO, Tasker RR, et al. Single-unit analysis of the human ventral thalamic nuclear group: somatosensory responses. *J Neurophysiol* 1988;59:299–316.
6. Schaltenbrand G, Wahren W. Atlas for stereotaxy of the human brain. Stuttgart: Thieme, 1977.
7. Allison T, McCarthy G, Wood CC, et al. Human cortical potentials evoked by stimulation of the median nerve. I. Cytoarchitectonic areas generating short-latency activity. *J Neurophysiol* 1989;62:694–710.

NeuroImages

Figure. (A) Normal EEG and ECG at beginning of registration. (B) Rhythmic activity of about 3/second in the right frontotemporal region accompanied by slight reduction of heart rate. (C) Spread of rhythmic activity to the left hemisphere; ECG shows asystole for 5 seconds.

Cardiac arrest during partial seizure

Peter Fuhr, MD, David Leppert, MD,
Basel, Switzerland

A 69-year-old man was admitted to the hospital after a brief confusional state following awakening, during which he fell. He had a history of mitral valve replacement 17 months earlier. On admission, he was alert and oriented, and neurologic findings were normal.

Results of EEG initially were normal and the ECG showed a sinus rhythm at 70 beats per minute (bpm) and a P-R interval of 0.13 seconds (figure, A). After a few minutes, theta waves appeared over the right frontotemporal region and became rhythmic, the heart rate fell to 52 bpm, and the P-R interval increased to 0.16 seconds (figure, B). Thirty seconds later, the rhythmic EEG activity spread contralaterally, the heart rate fell to 36 bpm, and the P-R

interval increased to 0.23 seconds, followed by asystole for 5 seconds (figure, C). At that time, the patient opened his eyes and smiled, but was unresponsive. Electric seizure activity stopped after 10 seconds. After 7 more seconds, the heart rate was 55 bpm, and a few minutes later, EEG and ECG had returned to normal. The patient was slightly disoriented immediately after the event but later on that day his mental status returned to normal.

Cardiac asystole induced by a complex partial seizure is a possible cause of sudden death in epilepsy, and primary cardiac arrhythmia causing syncope or presyncope is a likely misdiagnosis if cardiac monitoring is done without EEG.^{1,2}

1. Liedholm LJ, Gudjonsson O. Cardiac arrest due to partial epileptic seizure. *Neurology* 1992;42:824–829.
2. Jallon P. Arrhythmic seizures. *Epilepsia* 1997;38(suppl 11):S43–S47.

Neurology[®]

Cardiac arrest during partial seizure

Neurology 2000;54:2026

DOI 10.1212/WNL.54.10.2026

This information is current as of May 23, 2000

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/54/10/2026.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.