- hemolytic streptococcal infection. *J Child Neurol* 1998;13:354–356.
29. Muffat-Joly M, Barry B, Henin D, Fay M, Gehanno P, Pocard JJ. Orogenic meningoencephalitis induced by *Streptococcus pneumoniae* in gerbils. *Arch Otolaryngol Head Neck Surg* 1994;120:925–930.
 30. Blanot S, Muffat-Joly M, Vilde F, et al. A gerbil model for rhombencephalitis due to *Listeria monocytogenes*. *Microb Pathog* 1997;23:39–48.
 31. Watanabe-Ohnishi R, Aelion J, leGros L, et al. Characterization of unique human TCR V beta specificities for a family of streptococcal superantigens represented by rheumatogenic serotypes of M protein. *J Immunol* 1994;152:2066–2073.
 32. Toyosaki T, Yoshioka T, Tsuruta Y, Yutsudo T, Iwasaki M, Suzuki R. Definition of the mitogenic factor (MF) as a novel streptococcal superantigen that is different from streptococcal pyrogenic exotoxins A, B, and C. *Eur J Immunol* 1996;26:2693–2701.
 33. Rikiishi H, Okamoto S, Sugawara S, Tamura K, Liu ZX, Kumagai K. Superantigenicity of helper T cell mitogen (SPM-2) isolated from culture supernatants of streptococcus pyogenes. *Immunology* 1997;91:406–413.
 34. Vandevyver C, Mertens N, van den Elsen P, Medaer R, Raus J, Zhang J. Clonal expansion of myelin basic protein-reactive

- T cells in patients with multiple sclerosis: restricted T cell receptor V gene rearrangements and CDR3 sequence. *Eur J Immunol* 1995;25:958–968.
35. Maillard I, Luth F, Acha-Orbea H, Diggelman H. Role of the immune response induced by superantigens in the pathogenesis of microbial infections. *Parasitol* 1997;115:S67–S78.
 36. Herrmann T, Baschieri S, Less RK, MacDonald HR. In vivo responses of CD4+ and CD8+ cells to bacterial superantigens. *Eur J Immunol* 1992;22:1935–1938.
 37. Herrmann T, MacDonald HR. The CD8 T cell response to staphylococcal enterotoxins. *Semin Immunol* 1993;5:33–39.
 38. Hermans G, Denzer U, Lohse A, Raus J, Stinissen P. Cellular and humoral immune response against autoreactive T cells in multiple sclerosis patients after T cell vaccination. *J Autoimmun* 1999;13:233–246.
 39. Pohl-Koppe A, Burchett SK, Thiele EA, Hafler DA. Myelin basic protein reactive Th2 T cells are found in acute disseminated encephalomyelitis. *J Neuroimmunol* 1998;91:19–27.
 40. Genain CP, Abel K, Belmar N, et al. Late complications of immune deviation therapy in a nonhuman primate. *Science* 1996;274:2054–2057.
 41. Franchimont D, Louis E, Dewe W, et al. Effects of dexamethasone on the profile of cytokine secretion in human whole blood cell cultures. *Regul Pept* 1998;73:59–65.

NeuroImages

Figure. MRI at admission (postcontrast T1-weighted). (A) Ring-enhancing left cerebellar lesion and meningeal contrast enhancement. (B) Multiple intramedullary lesions.

Cerebellar and medullary histoplasmosis

M.J. Vos, MD, Y.J. Debets-Ossenkopp, MD, PhD,
F.A.P. Claessen, MD, G.J. Hazenberg, MD,
J.J. Heimans, MD, PhD, Amsterdam, the Netherlands

A 16-year-old Surinam boy presented with a 3-month history of progressive paraparesis and periods of fever and neck stiffness. Neurologic examination demonstrated a se-

vere spastic paraparesis, Babinski signs, and sensory disturbance under level T4. The patient was not taking immunosuppressants, was HIV-seronegative, and had normal CD4⁺ lymphocyte counts. Postcontrast T1-weighted MRI of brain and spinal cord revealed a ring-enhancing left cerebellar lesion, meningeal contrast enhancement, and multiple intramedullary lesions (figure). CSF examination showed a lymphocytic pleocytosis, elevated total protein, lowered glucose, and negative Gram staining and cultures. Biopsy of one intramedullary lesion disclosed granulomatous lymphohistiocytic inflammation with giant cells and necrosis; cultures were positive for *Histoplasma capsulatum*. Some neurologic improvement occurred after sequential treatment with amphotericin B and itraconazole.

Primary neurologic presentation of histoplasmosis is extremely rare, especially in young, nonimmunocompromised hosts. Tan et al. reported a case of disseminated histoplasmosis in a 72-year-old man, which was the first report of histoplasmosis presenting as a myelopathy.¹ Tabbal and Harik recently reported cerebral histoplasmosis in a 60-year-old man, presenting as a bilateral pyramidal syndrome.²

To our knowledge, this is the first case of cerebellar, meningeal, and medullary histoplasmosis reported in a young, nonimmunocompromised host. The severity of the neurologic presentation, lack of systemic manifestations, and extensiveness of CNS lesions on MRI are exceptional.

1. Tan V, Wilkins P, Badve S, et al. Histoplasmosis of the central nervous system. *JNNP* 1992;55:619–622.
2. Tabbal SD, Harik SI. Cerebral histoplasmosis. *N Engl J Med* 1999;340:1176.

Neurology®

Cerebellar and medullar histoplasmosis

Neurology 2000;54;1441

DOI 10.1212/WNL.54.7.1441

This information is current as of April 11, 2000

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/54/7/1441.full>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://n.neurology.org/content/54/7/1441.full##otherarticles>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.