20. Ivanova R, Henon N, Lepage V, et al. HLA-DR alleles display sex-dependent effects on survival and discriminate between individual and familial longevity. *Hum Mol Genet* 1998;7:187–194.
21. Woolf B. On estimating the relation between blood group and disease. *Ann Eugen* 1955;19:251–253.
22. Lewontin RC. The detection of linkage disequilibrium in molecular sequence data. *Genetics* 1995;140:377–388.
23. Payami H, Joe S, Farid NR, et al. Relative predispositional effects (RPEs) of marker alleles with disease: HLA-DR alleles and Graves disease. *Am J Hum Genet* 1989;45:541–546.
24. Spielman RS, McGinnis RE, Ewens WJ. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am J Hum Genet* 1993;52:506–516.
25. Kruglyak L, Daly MJ, Reeve-Daly MP, et al. Parametric and nonparametric linkage analysis: a unified multipoint approach. *Am J Hum Genet* 1996;58:1347–1363.
26. Sham PC, Curtis D. An extended transmission/disequilibrium test (TDT) for multi-allele marker loci. *Ann Hum Genet* 1995;59:323–336.
27. Hoch W, McConville J, Helms S, Newsom-Davis J, Melms A, Vincent A. Autoantibodies to the receptor tyrosine kinase MuSK in patients with myasthenia gravis without acetylcholine receptor antibodies. *Nat Med* 2001;7:365–368.
28. Kruglyak L, Lander E. Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nat Genet* 1995;11:241–247.
29. Degli-Esposti MA, Andreas A, Christiansen FT, et al. An approach to the localization of the susceptibility genes for generalized myasthenia gravis by mapping recombinant ancestral haplotypes. *Immunogenetics* 1992;35:355–364.
30. Janer M, Cowland A, Picard J, et al. A susceptibility region for myasthenia gravis extending into the HLA-class I sector telomeric to HLA-C. *Hum Immunol* 1999;60:909–917.
31. Matsuki K, Juji T, Tokunaga K, et al. HLA antigens in Japanese patients with myasthenia gravis. *J Clin Invest* 1990;86:392–399.
32. Muller-Hermelink HK, Marx A. Pathological aspects of malignant and benign thymic disorders. *Ann Med* 1999;31(suppl 2):5–14.
33. Spielman RS, Ewens WJ. A sibship test for linkage in the presence of association: the sib transmission/disequilibrium test. *Am J Hum Genet* 1998;62:450–458.

NeuroImages

Figure. (A) T1-weighted MRI; mixed pattern of CNS cryptococcus. (B) Virchow–Robin space crowded with cryptococci (hematoxylin and eosin, ×400 before reduction). Inset: cryptococcus neoformans with the mucinous capsule (hematoxylin and eosin ×1,000 before reduction). (C) T2-weighted MRI; mucoid material within Virchow–Robin spaces. Gelatinous pseudocysts in the head of the caudate and putamen.

Overwhelming CNS cryptococcus in AIDS

H.M. Bos, MD, P.A.M. Hofman, MD, PhD, G. Schreij, MD, PhD, F.J. Bot, MD, PhD, R.J. van Oostenbruggen, MD, PhD, Maastricht, the Netherlands

A 46-year-old HIV-positive man (CD₄ count 6 cells/mL, plasma HIV load 346×10^3 copies/mL) presented with a subacute onset of confusion. Neurologic examination demonstrated meningeal irritation. CT examination showed no abnormalities. CSF analysis revealed a cryptococcal infection. Because the patient deteriorated despite treatment, an MRI scan was performed 4 weeks later. The images showed a mixed pattern of dilated Virchow–Robin spaces filled with mucoid material,¹ gelatinous pseudocysts in the head of the caudate and putamen,² and widespread parenchymal and leptomeningeal nodules² (figure, A and C).

This is a classical example of a mixed pattern of CNS cryptococcus in a patient with AIDS. Histopathologic analysis showed a dilated Virchow–Robin space, crowded with cryptococci (hematoxylin and eosin, ×400; see the figure, B). Typically, no inflammation reaction and only limited gliosis in the surrounding nervous tissue was seen. The inset in figure B shows cryptococcus neoformans with the cell wall (dark) surrounded by a clear space representing the mucinous capsule (hematoxylin and eosin, ×1,000; see the figure, B). The patient died 2 months after receiving intensive antifungal therapy.

1. Mathews VP, Alo PL, Glass JD, Kumar AJ, McArthur JC. AIDS-related CNS cryptococcosis: radiologic–pathologic correlation. *AJNR Am J Neuroradiol* 1992;13:1477–1486.
2. Tien RD, Chu PK, Hesselink JR, Duberg A, Wiley C. Intracranial cryptococcosis in immunocompromised patients: CT and MR findings in 29 cases. *AJNR Am J Neuroradiol* 1991;12:283–289.

Neurology®

Overwhelming CNS cryptococcus in AIDS

H. M. Bos, P. A.M. Hofman, G. Schreij, et al.

Neurology 2001;57;1560

DOI 10.1212/WNL.57.9.1560

This information is current as of November 13, 2001

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/57/9/1560.full
References	This article cites 2 articles, 2 of which you can access for free at: http://n.neurology.org/content/57/9/1560.full#ref-list-1
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.