derly. Prevalence of Parkinson's disease and related disorders assessed by a door-to-door survey of inhabitants older than 65 years. *Arch Neurol* 1995;52:1017-1022.

12. Morgante L, Rocca WA, Di Rosa AE, et al. Prevalence of Parkinson's disease and other types of parkinsonism: a door-to-door survey in three Sicilian municipalities. The Sicilian Neuro-Epidemiologic Study (SNES) Group. *Neurology* 1992; 42:1901-1907.
13. Bharucha NE, Bharucha EP, Bharucha AE, Bhise AV, Schoenberg BS. Prevalence of Parkinson's disease in the Parsi community of Bombay, India. *Arch Neurol* 1988;45:1321-1323.
14. Wang SJ, Fuh JL, Teng EL, et al. A door-to-door survey of Parkinson's disease in a Chinese population in Kinmen. *Arch Neurol* 1996;53:66-71.
15. Li SC, Schoenberg BS, Wang CC, et al. A prevalence survey of Parkinson's disease and other movement disorders in the People's Republic of China. *Arch Neurol* 1985;42:655-657.
16. Menniti-Ippolito F, Spila-Alegiani S, Vanacore N, et al. Estimate of parkinsonism prevalence through drug prescription histories in the Province of Rome, Italy. *Acta Neurol Scand* 1995;92:49-54.
17. Mayeux R, Denaro J, Hemenegildo N, et al. A population-based investigation of Parkinson's disease with and without dementia. Relationship to age and gender. *Arch Neurol* 1992; 49:492-497.
18. D'Allessandro R, Gamberini G, Granieri E, Benassi G, Naccarato S, Manzaroli D. Prevalence of Parkinson's disease in the Republic of San Marino. *Neurology* 1987;37:1679-1682.
19. Wermuth L, Joensen P, Bungner N, Jeune B. High prevalence of Parkinson's disease in the Faroe Islands. *Neurology* 1997; 49:426-432.

20. Bennett DA, Beckett LA, Murray AM, et al. Prevalence of parkinsonian signs and associated mortality in a community population of older people. *N Engl J Med* 1996;334: 71-76.
21. Tandberg E, Larsen JP, Nessler EG, Riise T, Aarli JA. The epidemiology of Parkinson's disease in the county of Rogaland, Norway. *Mov Disord* 1995;10:541-549.
22. Baldereschi M, Di Carlo A, Rocca WA, et al. Parkinson's disease and parkinsonism in a longitudinal study: two-fold higher incidence in men. ILSA Working Group Italian Longitudinal Study on Aging. *Neurology* 2000;55:1358-1363.
23. Ben-Shlomo Y, Marmot MG. Survival and cause of death in a cohort of patients with parkinsonism: possible clues to aetiology. *J Neurol Neurosurg Psychiatry* 1995;58:293-299.
24. Miller DB, Ali SF, O'Callaghan JP, Laws SC. The impact of gender and estrogen on striatal dopaminergic neurotoxicity. *Ann NY Acad Sci* 1998;844:153-165.
25. Martilla RJ, Rinne UK. Epidemiology of Parkinson's disease in Finland. *Acta Neurol Scand* 1976;53:81-102.
26. Rosati G, Granieri E, Pinna L, et al. The risk of Parkinson's disease in Mediterranean people. *Neurology* 1980;30:250-255.
27. Okada K, Kobayashi S, Tsunematsu T. Prevalence of Parkinson's disease in Izumo City, Japan. *Gerontology* 1990;36:340-344.
28. Bower JH, Maraganore DM, McDonnell SK, Rocca WA. Incidence and distribution of parkinsonism in Olmsted County, Minnesota, 1976-1990. *Neurology* 1999;52:1214-1220.
29. de Rijk MC, Breteler MM, Graveland GA, et al. Prevalence of Parkinson's disease in the elderly: the Rotterdam Study. *Neurology* 1995;45:2143-2146.

Figure. Diffusion weighted imaging (echoplanar, repetition time = 5100 milliseconds, echo time = 137 milliseconds, b = 1000 T) showing a rim of high signal surrounding a lesion that does not demonstrate signal change or mass effect.

Progressive multifocal leukoencephalopathy

R.D. Henderson, FRACP, M.G. Smith, MBBS, P. Mowat, FRACP, and S.J. Read, MBBS, PhD, FRACP, Brisbane, Australia

A 58-year-old woman admitted to the hospital had a 3-month history of progressive neurologic deterioration

characterized by severe global aphasia and bilateral limb weakness, greater on the right. The patient had a 10-year history of systemic lupus erythematosus (SLE). Polyarthrititis had been present for 6 weeks before the onset of neurologic symptoms; she was treated with prednisone and methotrexate. A diagnosis of progressive multifocal leukoencephalopathy (PML) was established by a positive PCR for JC virus from CSF. The patient deteriorated and further active care was withdrawn.

Serial MRI showed an enlarging area of hypointensity on T1 and hyperintensity on T2-weighted images in the left parietal lobe. The lesion involved predominantly white matter and was without mass effect. A rim of high signal was present at the margins of the left parietal lesion on the diffusion-weighted image (figure) with a reduced apparent diffusion coefficient (ADC) on the ADC map. There was no enhancement after administration of IV gadolinium.

PML in SLE treated with immunosuppression is uncommon.¹ The MRI findings in PML often include circumscribed lesions predominantly in white matter without mass effect or contrast enhancement.^{1,2} The MR sequences indicate a progressing area of cytotoxic edema surrounding prior areas of white matter damage.²

1. Ahmed F, Aziz T, Kaufman LD. Progressive multifocal leukoencephalopathy in a patient with systemic lupus erythematosus. *J Rheumatol* 1999; 26:1609-1612.
2. Ohta K, Obara K, Sakauchi M, Obara K, Takane H, Yogo Y. Lesion identification by diffusion-weighted magnetic resonance imaging in progressive multifocal leukoencephalopathy. *J Neurol* 2001;248:809-811.

Neuro Images

Neurology®

Progressive multifocal leukoencephalopathy

R. D. Henderson, M. G. Smith, P. Mowat, et al.

Neurology 2002;58;1825

DOI 10.1212/WNL.58.12.1825

This information is current as of June 25, 2002

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/58/12/1825.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/58/12/1825.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.