Figure. High-resolution time-of-flight MR angiography (MRA) of the intracranial arteries (view from the top). (A) Multiple narrowing in several vascular territories (arrowheads). (B) A 71-day follow-up MRA shows regular vessels.

MR angiography in migrainous vasospasm

A. Schluter, MD, and B. Kissig, MD, Halle, Germany

A 60-year-old woman with a history of migraine without aura presented with severe headache attacks provoked by swimming in cold water lasting 1 to 6 hours accompanied by vomiting, nausea, and photophobia. The patient had no neurologic symptoms such as visual aura, hemiparesis, hemisensory loss, dysphasia, or vertigo related to the onset of the headache or the time of the first MR angiography (MRA). Initial self-medication with ergotamine and acetylsalicylic acid was ineffective. Blood pressure, blood indices,

and tests for vasculitis-associated antibodies were normal. On CT and MRI, we found no evidence for subarachnoid hemorrhage or cerebral edema. MRA done after 6 day-safer onset showed narrowing of multiple branches of cerebral arteries (figure, A). The headache attacks improved promptly to a combination of 1 mg ergotamine and 200 mg propyphenazone given subsequently to MRI. Clinical symptoms declined completely within 3 weeks. At 71-day follow-up MRA (figure, B) depicted normal vessels. The known evidence of vasospasm in prolonged migraine^{1,2} is most likely an epiphenomenon unrelated to headaches.² We cannot entirely rule out that initial administration of ergotamine did contribute to the vessel situation.

Address correspondence and reprint requests to Dr. A. Schluter, Martin-Luther-University Halle-Wittenberg, Department of Radiology, Ernst-Grube-Strasse 40, 06097 Halle, Germany.

1. Rothrock JF, Walicke P, Swenson MR, Lyden PD, Logan WR. Migrainous stroke. *Arch Neurol* 1988;45:63–67.
2. Solomon S, Lipton RB, Harris PY. Arterial stenosis in migraine: spasm or arteriopathy? *Headache* 1990;30:52–61.

Neurology[®]

MR angiography in migrainous vasospasm

A. Schluter and B. Kissig

Neurology 2002;59;1772

DOI 10.1212/01.WNL.0000033278.68801.22

This information is current as of December 10, 2002

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/59/11/1772.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/59/11/1772.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/59/11/1772.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Migraine http://n.neurology.org/cgi/collection/migraine MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.