References

1. Becker G, Berg D, Rausch W-D, Lange HKW, Riederer P, Reiners K. Increased tissue copper and manganese content in primary adult-onset dystonia. *Ann Neurol* 1999;46:260–263.
2. Berg D, Weishaupt A, Francis MJ, et al. Changes of copper-transporting proteins and ceruloplasmin in the lentiform nuclei in primary adult-onset dystonia. *Ann Neurol* 2000;47:827–830.
3. Kruse N, Berg D, Francis MJ, et al. Reduction of Menkes mRNA and copper in leukocytes of patients with primary adult-onset dystonia. *Ann Neurol* 2001;49:405–408.
4. Parker N. Hereditary whispering dysphonia. *J Neurol Neurosurg Psychiatry* 1985;48:218–224.
5. Tümer Z, Lund C, Tolshave J, Vural B, Tonnesen T, Horn N. Identification of point mutations in 41 unrelated patients affected with Menkes disease. *Am J Hum Genet* 1997;60:63–71.
6. Petrushkin K, Lutsenko S, Chernov I, Ross BM, Kaplan JH, Gilliam TC. Characterization of the Wilson disease gene encoding a P-type copper transporting ATPase: genomic organization, alternative splicing, and structure/function predictions. *Hum Mol Genet* 1994;3:1647–1656.
7. Das S, Levinson B, Whitney S, Vulpe C, Packman S, Gitschier J. Diverse mutations in patients with Menkes disease often lead to exon skipping. *Am J Hum Genet* 1994;55:883–889.

Pure sensory stroke from cortical infarction

David A. Wolk, MD, Steven R. Messé, MD, Scott E. Kasner, MD, Philadelphia, PA

A 52-year-old man developed sudden numbness and paresthesias of the left face, ear, arm, leg, and trunk, but not the tongue or genitals, equally involving all affected areas. All elementary modalities of pain/temperature, light touch, and proprioception were impaired. He also had mild left-sided heaviness. On examination, he had normal visual fields, normal strength, and no neglect or visuospatial deficits. MRI showed infarction in the right postcentral gyrus (figure). The etiology of the stroke was cryptogenic. Over the next few months the symptoms in his leg improved, but he gradually developed chronic left arm dysesthesia, similar to the Roussy-Déjérine syndrome associated with thalamic strokes. He later developed tactile hallucinations in his left hand.

The anatomic localization of sensory loss in stroke generally is reliably determined by evaluation of the distribution of the deficit and the sensory modalities involved, but this can be misleading in the acute setting. The putative dictum is that in the absence of neuropsychological dysfunction, a pure sensory stroke involving face, arm, leg, and trunk usually localizes to the thalamus,¹ or rarely to the brainstem. Involvement of two or more regions of the body has high positive predictive value (95%) for a small vessel ischemic event.² Rarely, parietal cortical infarction may cause a pseudo-thalamic syndrome, particularly in

Figure. MRI FLAIR sequences revealed increased signal intensity isolated to the postcentral gyrus. The infarction was limited to the gray matter and was extended approximately 2 cm above and below this representative slice.

the acute setting. Over time the clinical manifestations may become more suggestive of cortical localization.

Address correspondence and reprint requests to Dr. Scott E. Kasner, Dept. of Neurology, Comprehensive Stroke Center, University of Pennsylvania Medical Center, 3W Gates Bldg., 3400 Spruce St., Philadelphia, PA 19104; e-mail: Kasner@mail.med.upenn.edu

1. Bassetti C, Bogousslavsky J, Regli F. Sensory syndromes in parietal stroke. *Neurology* 1993;43:1942–1949.
2. Chimowitz MI, Furlan AJ, Sila CA, Parandhi L, Beck GJ. Etiology of motor or sensory stroke: a prospective study of predictive value of clinical and radiographic features. *Ann Neurol* 1991;30:519–525.

Neurology®

Pure sensory stroke from cortical infarction

David A. Wolk, Steven R. Messé and Scott E. Kasner

Neurology 2002;59:783

DOI 10.1212/WNL.59.5.783

This information is current as of September 10, 2002

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/59/5/783.full
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/59/5/783.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Infarction http://n.neurology.org/cgi/collection/infarction MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.