Figure. T2-weighted fluid-attenuated inversion recovery (A) and spin-echo (B) MR images at the level of the pons reveal a dolichoectatic basilar artery aneurysm and a thrombus partially filling the lumen (clear arrow). A probable stroke in the pons involving the pontine perforating artery territory is also present (A and B, solid arrow). Thrombus was not present 3 years ago (clear arrow in C).

Basilar artery aneurysm thrombosis

Pitchaiah Mandava, MD, PhD; Ann Bebensee, MD;
R. Ryan Rawlings, DO; and Thomas A. Kent, MD,
Galveston, TX

A 53-year-old man presented with new-onset severe right-sided weakness, dysarthria, dysphagia, and nystagmus. Posterior circulation infarct was suspected and MRI showed a thrombus in a previously characterized dolichoectatic basilar artery aneurysm (figure). The evolution of an arterial thrombus by MRI differs from an intraparenchymal hematoma.¹ Hyperintensity on T1 (not shown) and

T2 spin-echo images indicate an arterial thrombus hours to days old and may reflect accumulation of paramagnetic methemoglobin and changes in protein hydration as red blood cells lyse.¹

This and two prior strokes were preceded by an acute gout attack. Infection and inflammatory conditions including gout may be triggers for ischemic events,² conceivably increasing the risk for thrombus formation at a vulnerable site.

Address correspondence and reprint requests to Dr. Pitchaiah Mandava, Department of Neurology, The University of Texas Medical Branch, 301 University Blvd., Galveston, TX 77555-0539; e-mail: pmandava@utmb.edu

1. Corti R, Osende JJ, Fayad ZA, et al. In vivo noninvasive detection and age definition of arterial thrombus by MRI. *J Am Coll Cardiol* 2002;39:1366–1373.
2. Macko RF, Ameriso SF, Barndt R, Clough W, Weiner JM, Fisher M. Precipitants of brain infarction. Roles of preceding infection/inflammation and recent psychological stress. *Stroke* 1996;27:1999–2004.

Neurology®

Basilar artery aneurysm thrombosis

Pitchaiah Mandava, Ann Bebensee, R. Ryan Rawlings, et al.

Neurology 2002;59;1287

DOI 10.1212/01.WNL.0000033623.85661.92

This information is current as of October 22, 2002

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/59/8/1287.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/59/8/1287.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
All Cerebrovascular disease/Stroke
http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke
All Imaging
http://n.neurology.org/cgi/collection/all_imaging
Infarction
<http://n.neurology.org/cgi/collection/infarction>
MRI
<http://n.neurology.org/cgi/collection/mri>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.