28. Call DR, Nemzek JA, Ebong SJ, Bolgos GL, Newcomb DE, Remick DG. Ratio of local to systemic chemokine concentrations regulates neutrophil recruitment. *Am J Pathol* 2001;158:715–721.
29. Fazioli F, Resnati M, Sidenius N, Higashimoto Y, Appella E, Blasi F. A urokinase-sensitive region of the human urokinase receptor is responsible for its chemotactic activity. *EMBO J* 1997;16:7279–7286.
30. Winkler F, Koedel U, Kastenbauer S, Pfister HW. Differential expression of nitric oxide synthases in bacterial meningitis: role of the inducible isoform for blood-brain barrier breakdown. *J Infect Dis* 2001;183:1749–1759.
31. Kastenbauer S, Koedel U, Becker BF, Pfister HW. Oxidative stress in bacterial meningitis in humans. *Neurology* 2002;58:186–191.
32. Pfister HW, Scheld WM. Brain injury in bacterial meningitis: therapeutic implications. *Curr Opin Neurol* 1997;10:254–259.
33. Kataoka K, Asai T, Taneda M, et al. Roles of urokinase type plasminogen activator in a brain stab wound. *Brain Res* 2000;887:187–190.
34. Philippe J, Offner F, Declerck PJ, et al. Fibrinolysis and coagulation in patients with infectious disease and sepsis. *Thromb Haemost* 1991;65:291–295.
35. Philippe J, Dooijewaard G, Offner F, Turion P, Baele G, Leroux-Roels G. Granulocyte elastase, tumor necrosis factor- α and urokinase levels as prognostic markers in severe infection. *Thromb Haemost* 1992;68:19–23.
36. Robbie LA, Dummer S, Booth NA, Adey GD, Bennett B. Plasminogen activator inhibitor 2 and urokinase-type plasminogen activator in plasma and leucocytes in patients with severe sepsis. *Br J Haematol* 2000;109:342–348.
37. Creange A, Sharshar T, Planchenault T, et al. Matrix metalloproteinase-9 is increased and correlates with severity in Guillain-Barré syndrome. *Neurology* 1999;53:1683–1691.

Neuro Images

Figure. Cells in cultured 200- μ m-thick motor cortex slices from a deceased PD patient (female, 79 years, postmortem delay: 5 hours) express the reporter genes β -galactosidase (A) and green fluorescent protein (B). The slices were infected with recombinant adeno-associated virus containing the respective genes after 34 (A) and 20 (B) days in culture and stained 10 (A) and 12 (B) days later. (A), pyramidal cells from layer III are shown. (B) An unidentified cell located in layer V, which may be an interneuron or an astrocyte. Bar denotes 20 μ m.

Life after death?

R.W.H. Verwer, PhD, W.T.J.M.C. Hermens, PhD, O. ter Brake, J. Verhaagen, PhD, and D.F. Swaab, MD, PhD, Amsterdam, the Netherlands

It is generally accepted that irreparable damage to the brain follows if its blood supply is interrupted for more than 5 to 10 minutes.¹ This is certainly true for the brain,

functioning as an integral organ. However, we obtained evidence that many neurons survive a postmortem delay of 2 to 8 hours and can be maintained alive outside the body for extensive periods of time.² Thus, it appears that neurons are less vulnerable to hypoxia/ischemia than is often assumed. Figures A and B illustrate that neurons and other cells in long-term slice cultures from postmortem human brain can still express viral-vector-mediated transgenes for β -galactosidase and green fluorescent protein. This shows that neurons may be amenable to gene therapeutic approaches after ischemic insults.

The authors are unaware of any conflict of interest.

Supported by Internationale Stichting Alzheimer Onderzoek (ISAO), Alzheimer Stichting Nederland, Platform voor Alternatieven voor Dierproeven, New Drug Research Foundation, Research Institute for Diseases in the Elderly (RIDE/NOW), and Netherlands Organisation for Scientific Research (NOW/GB-MW). Brain material was provided by the Netherlands Brain Bank (coordinator: Dr. R. Ravid).

Address correspondence and reprint requests to Dr. R.W.H. Verwer, Netherlands Institute for Brain Research, Meibergdreef 33, 1105AZ Amsterdam, the Netherlands; e-mail: r.verwer@nih.knaw.nl

1. van Lommel P, van Wees R, Meyers V, Elfferich I. Near-death experience in survivors of cardiac arrest: a prospective study in the Netherlands. *Lancet* 2001;358:2039–2045.
2. Verwer RWH, Hermens WTJMC, Dijkhuizen PA, et al. Cells in human postmortem brain tissue slices remain alive for several weeks in culture. *FASEB J* 2002;16:54–60.

Neurology[®]

Life after death?

R.W.H. Verwer, W.T.J.M.C. Hermens, O. ter Brake, et al.

Neurology 2002;59;1355

DOI 10.1212/01.WNL.0000032758.08215.D1

This information is current as of November 12, 2002

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/59/9/1355.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/59/9/1355.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
All Cerebrovascular disease/Stroke
http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke
Infarction
<http://n.neurology.org/cgi/collection/infarction>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.