Figure. (A) T1-weighted sagittal image shows hyperintense signal along the left corticospinal tract. This striking appearance is also present on the opposite hemisphere. (B) Axial FLAIR image shows hyperintensity of precentral gyrus bilaterally. (C) Axial FLAIR image shows hyperintense corticospinal tract at the crus cerebri. This series of MR images clearly demonstrated bilateral corticospinal tract involvement in amyotrophic lateral sclerosis.

MRI in ALS: Corticospinal tract hyperintensity

Yong Chern Lee, MBBS, Romesh Markus, MBChB, FRACP, and Andrew Hughes, MD, FRACP, Heidelberg, Victoria, Australia

A 32-year-old woman with no family history presented with 12 months of progressive left upper limb weakness. Examination revealed moderate wasting and pyramidal weakness in all muscle groups in the left upper limb, especially the shoulder girdle and

intrinsic hand muscles. Mild pyramidal weakness was also noted in the left lower limb. No fasciculation was noted. She was hyper-tonic and hyperreflexic in all four limbs. Clonus and upgoing plan-tar response were elicited bilaterally. Sensory and cranial nerve examinations were unremarkable. Neurophysiology and MRI studies (figure) supported the diagnosis of amyotrophic lateral sclerosis (ALS). Recent neuroimaging studies have demonstrated hyperintense signals on T2-weighted and fluid-attenuated inversion recovery images along the corticospinal tract in some patients with ALS.¹ This is thought to be due to a dying back process.

Address correspondence and reprint requests to Dr. Yong Chern Lee, Neurology Department, 3KZ Building, Austin and Repatriation Medical Centre, Studley Road, Heidelberg 3084, Melbourne, Victoria, Australia; e-mail: yongchern@yahoo.com

1. Hecht MJ, Fellner F, Fellner C, Hilz MJ, Heuss D, Neundorfer B. MRI-FLAIR images of the head show corticospinal tract alterations in ALS patients more frequently than T2-, T1- and proton-density-weighted images. *J Neurol Sci* 2001;186:37–44.

Neurology®

MRI in ALS: Corticospinal tract hyperintensity
Yong Chern Lee, Romesh Markus and Andrew Hughes
Neurology 2003;61;1600
DOI 10.1212/01.WNL.0000096015.48322.2A

This information is current as of December 8, 2003

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/61/11/1600.full
References	This article cites 1 articles, 0 of which you can access for free at: http://n.neurology.org/content/61/11/1600.full#ref-list-1
Citations	This article has been cited by 3 HighWire-hosted articles: http://n.neurology.org/content/61/11/1600.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Imaging http://n.neurology.org/cgi/collection/all_imaging Amyotrophic lateral sclerosis http://n.neurology.org/cgi/collection/amyotrophic_lateral_sclerosis_MRI MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.