HSV encephalitis with posterior localization

Kouichi Ohta, MD; Mitsuhsa Tabata, MD; Ichiro Nakachi, MD; and Katsuyuki Obara, MD

A 50-year-old woman presented with generalized seizure, stupor, and fever. Lumbar puncture yielded CSF with increased leukocytes and protein content. PCR analyses of the CSF revealed DNA of only herpes simplex virus (HSV) type 1 among herpes viruses. Brain diffusion-weighted MR image clearly revealed multifocal hyperintense lesions (figure).¹ This is the first reported case of HSV encephalitis in which MRI demonstrated lesions restricted to the temporal, parietal, and occipital lobes without the typical localization of HSV encephalitis in the insula and inferior frontal lobe.² IV acyclovir proved effective to limit the extension of the lesions, resulting in her clinical recovery.

1. Ohta K, Funaki M, Tanaka M, Suzuki N. Early cerebellar involvement on diffusion-weighted magnetic resonance images in herpes simplex encephalitis. *J Neurol* 1999;246:736–738.
2. Chaudhuri A, Kennedy PG. Diagnosis and treatment of viral encephalitis. *Postgrad Med J* 2002;78:575–583.

From the Department of Neurology, Mito Red Cross Hospital, Japan.

Received March 5, 2003. Accepted June 15, 2003.

Address correspondence and reprints requests to Dr. K. Ohta, Department of Neurology, Mito Red Cross Hospital, 3-12-48 San-no-maru, Mito, 310-0011 Japan; e-mail: k-ohta@seagreen.ocn.ne.jp

Figure. Diffusion-weighted MR images of the brain on the 9th day of illness. Multiple hyperintense areas were identified in bilateral occipital lobes and left parietal lobe in addition to bilateral inferomedial temporal lobes. Those areas were also hyperintense on T2-weighted images, although less recognizable than on diffusion-weighted images.

DISAGREE? AGREE? HAVE A QUESTION? HAVE AN ANSWER?

Respond to an article in *Neurology* through our online Correspondence system:

- Visit www.neurology.org
- Access specific article on which you would like to comment
- Click on "Correspondence: Submit a response" in the content box
- Enter contact information
- Upload your Correspondence
- Press Send Response

Correspondence will then be transmitted to the *Neurology* Editorial Office for review. Accepted material will be posted within 10–14 days of acceptance. Selected correspondence will subsequently appear in the print Journal. See our Information for Authors at www.neurology.org for format requirements.

Neurology®

HSV encephalitis with posterior localization

Kouichi Ohta, Mitsuhsa Tabata, Ichiro Nakachi, et al.

Neurology 2003;61;862

DOI 10.1212/01.WNL.0000085870.71214.CA

This information is current as of September 22, 2003

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/61/6/862.full
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/61/6/862.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Encephalitis http://n.neurology.org/cgi/collection/encephalitis MRI http://n.neurology.org/cgi/collection/mri Viral infections http://n.neurology.org/cgi/collection/viral_infections
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.