Recurrent longitudinal myelitis as primary manifestation of SLE

F.-G. Lehnhardt, MD; P. Impekoven, MD; A. Rubbert, MD; L. Burghaus, MD; M. Neveling, MD; W.-D. Heiss, MD; and A.H. Jacobs MD, Köln, Germany

A 19-year-old woman presented with fever, neck pain, weakness of both legs, urinary retention, nuchal rigidity, quadriplegia, and a T6 sensory level. MRI showed hyperintensities (T2-weighted) of the entire spinal cord (figure), consistent with acute longitudinal myelitis (ALM). Laboratory studies confirmed the diagnosis of systemic lupus erythematosus (SLE) with antiphospholipid antibody syndrome (APS). A complete remission occurred after treatment with methylprednisolone and IV immunoglobulins. Symptoms recurred 13 months later; treatment was repeated, followed by cyclophosphamide.

The *continuous* involvement of spinal cord segments, or ALM, may be related to APS^{1,2} and distinct² from acute transverse myelitis (ATM).¹ Patients with ATM or ALM should be tested for SLE and APS, with treatment to include early IV corticosteroids followed by cyclophosphamide.

1. Kovacs B, Lafferty TL, Brent LH, DeHoratius RJ. Transverse myelopathy in systemic lupus erythematosus: an analysis of 14 cases and review of the literature. *Ann Rheum Dis* 2000;59:120–124.
2. Deodhar AA, Hochenadel T, Bennett RM. Longitudinal involvement of the spinal cord in a patient with lupus related transverse myelitis. *J Rheumatol* 1999;26:446–449.

Address correspondence and reprint requests to Priv.-Doz. Dr. A.H. Jacobs, Department of Neurology and MPI for Neurologic Research, Gleuelerstr. 50, 50931 Köln, Germany, e-mail: Andreas.Jacobs@pet.mpin-koeln.mpg.de

Figure. MRI of the spinal cord after first admission (A) and during recurrent episode 13 months later (B). T2-weighted sagittal images reveal an abnormal intramedullary high signal intensity and enlargement of the spinal cord reaching from the upper cervical segments to the conus medullaris.

Neurology®

Recurrent longitudinal myelitis as primary manifestation of SLE

F.-G. Lehnhardt, P. Impekoven, A. Rubbert, et al.

Neurology 2004;63;1976

DOI 10.1212/01.WNL.0000140623.47437.B3

This information is current as of November 22, 2004

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/63/10/1976.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/63/10/1976.full#ref-list-1>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://n.neurology.org/content/63/10/1976.full##otherarticles>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All Immunology

http://n.neurology.org/cgi/collection/all_immunology

All Spinal Cord

http://n.neurology.org/cgi/collection/all_spinal_cord

Lupus

<http://n.neurology.org/cgi/collection/lupus>

Transverse myelitis

http://n.neurology.org/cgi/collection/transverse_myelitis

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.