Figure. Cervical MRI upon admission: central spinal cord signal abnormalities, hyperintense in T2-weighted (A) and hypointense in T1-weighted and in fluid-attenuated inversion recovery (FLAIR) images (B), compatible with a hydromyelia. It coexisted with T2-weighted hyperintensities of the posterior, lateral, and anterior columns and completely remitted with vitamin B12 treatment.

Reversible cervical hydromyelia in subacute combined degeneration

N. Canas, MD, MS; S. Calado, MD; C. Ribeiro, MD; and J. Vale, MD, Lisbon, Portugal

A 53-year-old woman presented with insidiously progressive tetraparesis, C5 sensitive level and urinary incontinence for 16 months. Cervical MRI disclosed signal abnormalities consistent with subacute combined degeneration¹ (SCD), associated with a hydromyelia (figure). Laboratory studies confirmed the diagnosis

Address correspondence and reprint requests to Dr. Nuno Canas, Serviço Universitário de Neurologia, Hospital de Egas Moniz, Rua da Junqueira 126, 1349-019 Lisboa, Portugal; e-mail: nunocanas@netcabo.pt

of SCD. Clinical and imagiologic remission occurred with vitamin B12 treatment.

The MRI follow-up studies and the hydromyelia topography suggest that it was formed as a functional consequence of long-tract pathway demyelination.

The presence of hydromyelia, when associated with the characteristic MRI, clinical and laboratorial features of SCD, may be attributed to vitamin B12 deficiency itself.²

1. Pittock SJ, Payne TA, Harper MC. Reversible myelopathy in a 34-year-old man with vitamin B12 deficiency. *Mayo Clin Proc* 2002;77:291–294.
2. Puri V, Chaudhry N, Gulati P. Syringomyelia-like manifestation of subacute combined degeneration. *J Clin Neurosci* 2004;11:672–675.

Neurology®

Reversible cervical hydromyelia in subacute combined degeneration

N. Canas, S. Calado, C. Ribeiro, et al.

Neurology 2005;65;E7

DOI 10.1212/01.wnl.0000177925.92714.a9

This information is current as of August 8, 2005

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/65/3/E7.full>

Supplementary Material

Supplementary material can be found at:
<http://n.neurology.org/content/suppl/2007/04/02/65.3.E7.DC1>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/65/3/E7.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All Spinal Cord

http://n.neurology.org/cgi/collection/all_spinal_cord

Hematologic

<http://n.neurology.org/cgi/collection/hematologic>

Nutritional

<http://n.neurology.org/cgi/collection/nutritional>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.