VIDEO

Psychogenic paroxysmal dyskinesia following paroxysmal hemidystonia in multiple sclerosis

John C. Morgan, MD, PhD; Mary Hughes, MD;
Ramon E. Figueroa, MD; and Kapil D. Sethi, MD, FRCP,
Augusta, GA

A 27-year-old woman presented because of multiple daily episodes of painful “drawing” in her right arm and leg precipitated by hyperventilation (see video, segment 1). These episodes would typically last between 30 seconds and 2 minutes. Brain MRI and CSF analysis were consistent with multiple sclerosis (MS). MS-related paroxysmal hemidystonia was diagnosed because of a typical pattern of limb involvement, evolution of right hemibody dystonic posturing over seconds, the trigger (hyperventilation in the setting of MS), duration and frequency of dystonic spasms, and resolution of the spasms following acetazolamide treatment.¹ She subsequently began β -interferon injections for MS.

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. John C. Morgan, Medical College of Georgia, 1429 Harper Street, HF1121A, Augusta, GA; e-mail: jmorgan@mail.mcg.edu

One year later, the patient presented with “spasms” lasting up to 15 minutes (see video, segment 2) after missing a single β -interferon injection. Psychogenic paroxysmal dyskinesias were diagnosed because of an obvious trigger (concern over missing a single β -interferon injection), the prolonged evolution and duration of her “spasms,” bilateral representation and opisthotonic posturing during the “spasm” (a typical finding in psychogenic paroxysmal dyskinesias),² and her response to both suggestion and placebo. The episode recorded in segment 2 was typical of these events seen during this timeframe without induction.

Clinicians should have suspicion when patients present with an unusual paroxysmal dyskinesia, even in the setting of a previously documented “organic” dystonia.³

Copyright © 2005 by AAN Enterprises, Inc.

1. Sethi KD, Hess DC, Huffnagle VH, Adams RJ. Acetazolamide treatment of paroxysmal dystonia in central demyelinating disease. *Neurology* 1992;42:919–921.
2. Bruno MK, Hallett M, Gwinn-Hardy K, et al. Clinical evaluation of idiopathic paroxysmal kinesigenic dyskinesia: new diagnostic criteria. *Neurology* 2004;63:2280–2287.
3. Factor SA, Podskalny GD, Molho ES. Psychogenic movement disorders: frequency, clinical profile, and characteristics. *J Neurol Neurosurg Psychiatry* 1995;59:406–412.

Neurology[®]

Psychogenic paroxysmal dyskinesia following paroxysmal hemidystonia in multiple sclerosis

John C. Morgan, Mary Hughes, Ramon E. Figueroa, et al.

Neurology 2005;65;E12

DOI 10.1212/01.wnl.0000170367.33505.e9

This information is current as of September 26, 2005

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/65/6/E12.full>

Supplementary Material

Supplementary material can be found at:
<http://n.neurology.org/content/suppl/2005/09/23/65.6.E12.DC1>

References

This article cites 3 articles, 3 of which you can access for free at:
<http://n.neurology.org/content/65/6/E12.full#ref-list-1>

Citations

This article has been cited by 3 HighWire-hosted articles:
<http://n.neurology.org/content/65/6/E12.full##otherarticles>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.