Clinical findings of the phakomatoses: Sturge-Weber syndrome

Mark Quigg, MD, MSc; Robert S. Rust, MD; and James Q. Miller, MD†

Figure 1. (A) Ophthalmic port-wine stain (PWS) in an infant. Glaucoma is a frequent complication. PWS that spares ophthalmic trigeminal distribution is usually not indicative of underlying Sturge-Weber syndrome.¹ (B) Bilateral or more lower facial involvement occurs in 15%.¹ As in this child, about half have an extracranial nevus that typically respects segmental anatomy.

Figure 2. (A) A-P skull film that demonstrates the gyral calcifications of the subarachnoid angiomas of Sturge-Weber syndrome, tortuous parallel “tram tracks” across most of the right hemisphere. (B) In another patient, uncontrasted axial CT and (C) gradient-echo axial MRI show gyral calcifications, atrophy, and gliosis of the underlying cortex due to leptomeningeal involvement.

The dermatologic hallmark of Sturge-Weber syndrome (SWS) is the nevus flammeus (port-wine stain, PWS), a angiomatous lesion usually distributed in the region innervated by the ophthalmic branch of the trigeminal nerve¹ caused by lack of normal regression of embryonic vascular plexus of the cephalic neural tube. The

variable constellation of clinical findings of hemiparesis, mental retardation, and epilepsy is caused by the resulting leptomeningeal and cortical angiomatosis and calcification. SWS is a congenital and nonhereditary disorder. Physical findings of SWS are shown in figure 1 and neuroimaging in figure 2.

Acknowledgment

The authors thank C. Douglas Phillips of the Division of Neuroradiology for his contributions of MRI images.

Reference

1. Tallman B, Tan OT, Morelli JG, et al. Location of port-wine stains and the likelihood of ophthalmic and/or central nervous system complications. *Pediatrics* 1991;87:323–327.

From the Department of Neurology, University of Virginia, Charlottesville.

†Deceased.

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Mark Quigg, Department of Neurology, Box 800394, Health Sciences Center, University of Virginia, Charlottesville, VA 22908; e-mail: quigg@virginia.edu

Neurology®

Clinical findings of the phakomatoses: Sturge–Weber syndrome

Mark Quigg, Robert S. Rust and James Q. Miller

Neurology 2006;66;E17-E18

DOI 10.1212/01.wnl.0000197789.72780.a9

This information is current as of February 27, 2006

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/66/4/E17.full
References	This article cites 1 articles, 0 of which you can access for free at: http://n.neurology.org/content/66/4/E17.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Epilepsy/Seizures http://n.neurology.org/cgi/collection/all_epilepsy_seizures All Imaging http://n.neurology.org/cgi/collection/all_imaging MRI http://n.neurology.org/cgi/collection/mri Other neurocutaneous disorders http://n.neurology.org/cgi/collection/other_neurocutaneous_disorders
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.