Horner syndrome due to internal carotid artery dissection

Sara Mazzucco, MD; and Nicolò Rizzuto, MD

A few hours after a motorcycle accident with multiple rib and right wrist fractures, a 37-year-old man developed transient dysphasia and right hemiparesis associated with left Horner syndrome (figure A). Brain CT scan was normal, but diffusion weighted (DW) MRI disclosed recent ischemia (figure B). Occlusive dissection of left extracranial internal carotid artery (ICA) was demonstrated (figure C–E), and IV heparin infusion was started with a subsequent neurologically uneventful clinical course.

ICA dissection can occur even with unapparent neck injuries¹; therefore, trauma associated with Horner syndrome should instigate a workup for carotid dissection.²

Figure. (A) Horner syndrome (miosis and ptosis). (B) Multiple left frontoparietal acute ischemic lesions on DW MRI. (C) Carotid angiography with evidence of left extracranial ICA occlusion. (D) ICA mural hematoma due to dissection evident on MRI scan. (E) Intimal flap (arrows) appearance on Duplex scan, pathognomonic sign of dissection.

References

1. Miller PR, Fabian TC, Croce MA, et al. Prospective screening for blunt cerebrovascular injuries: analysis of diagnostic modalities and outcomes. *Ann Surg* 2002;236:386–395.
2. Chan CC, Paine M, O'Day J. Carotid dissection: a common cause of Horner's syndrome. *Clin Experiment Ophthalmol* 2001;29:411–415.

From the Department of Neurological Sciences and Vision, Section of Clinical Neurology, University of Verona, Italy.

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Sara Mazzucco, Clinica Neurologica, Policlinico G.B.Rossi, Piazzale L.A.Scuro,10, 37134 Verona, Italy; e-mail: sara.mazzucco@univr.it

Neurology[®]

Teaching NeuroImage: Horner syndrome due to internal carotid artery dissection

Sara Mazzucco and Nicolò Rizzuto

Neurology 2006;66;E19

DOI 10.1212/01.wnl.0000200954.21276.96

This information is current as of March 13, 2006

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/66/5/E19.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/66/5/E19.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Autonomic diseases http://n.neurology.org/cgi/collection/autonomic_diseases Carotid artery dissection http://n.neurology.org/cgi/collection/carotid_artery_dissection
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.