Figure 1. (A) The patient's own drawing of macropsia demonstrates disproportionate magnification of right face of a person. Numerous dots represent magnified hair pores. (B) The patient's own drawing shows the list of unformed hallucinations during seizure.

Figure 2. (A) Gradient echo imaging demonstrates multiple cavernous hemangiomas. (B) EEG discovered electrographic seizure in left occipital region.

Visual illusory and hallucinatory phenomena in a patient with left occipital seizures

Makoto Kawai, MD; Igor M. Chermes, MD; and
Ian L. Goldsmith, MD, Houston, TX

A 60-year-old woman had sudden onset transient visual symptoms in her right visual field after having worsening postural headaches for 1 year. Visual symptoms included

palinopsia or abnormally recurring visual imagery, macropsia, unformed hallucinations, and hemianopia (figure 1). The symptoms lasted from 5 minutes to a few hours. Imaging demonstrated multiple cavernous hemangiomas including one involving the left temporo-occipital region (figure 2). Electrographic left occipital seizures were correlated with unformed hallucinations. The symptoms resolved with topiramate.

Seizures in the occipital region are reported to cause unformed hallucinations or hemianopia, and rarely the ones involving temporo-parieto-occipital junction can present visual illusory symptoms.^{1,2}

Supported by Peter Kellaway, PhD, Endowment for Research, Baylor College of Medicine, Houston, TX.

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Makoto Kawai, Peter Kellaway Section of Neurophysiology, Department of Neurology, Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030; e-mail: makotokawai71@aol.com

1. Norton JW, Corbett JJ. Visual perceptual abnormalities: hallucinations and illusions. *Semin Neurol* 2000;20:111-121.
2. Blume WT, Wiebe S, Tapsell LM. Occipital epilepsy: lateral versus mesial. *Brain* 2005;128(Pt 5):1209-1225.

Neurology®

Visual illusory and hallucinatory phenomena in a patient with left occipital seizures

Makoto Kawai, Igor M. Chermes and Ian L. Goldsmith

Neurology 2006;67:1457

DOI 10.1212/01.wnl.0000227180.18972.e6

This information is current as of October 23, 2006

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/67/8/1457.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/67/8/1457.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All Epilepsy/Seizures

http://n.neurology.org/cgi/collection/all_epilepsy_seizures

Epilepsy semiology

http://n.neurology.org/cgi/collection/epilepsy_semiology

Hallucinations

<http://n.neurology.org/cgi/collection/hallucinations>

Partial seizures

http://n.neurology.org/cgi/collection/partial_seizures

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.