Teaching Video

Is it III alone, or III and IV?

Stephen G. Reich, MD

VIDEO

Address correspondence and reprint requests to Dr. Stephen G. Reich, University of Maryland School of Medicine, 22 South Greene Street, N4W46, Baltimore, MD 21201
sreich@som.umaryland.edu

The most important questions when confronted with an oculomotor (III) palsy are 1) Is the pupil spared? 2) Is it complete aside from pupil sparing? and 3) Is it in isolation? A “no” answer to any makes a benign, ischemic III palsy less likely.¹ In the presence of a III palsy, the traditional method of testing the trochlear nerve (IV) at the bedside by asking the patient to depress the adducted eye cannot be performed. Instead, the patient should be instructed to abduct the eye and then look down; if IV is intact, there will be intorsion.² Confirming that IV is intact in the presence of a III palsy is important because the combination of an oculomotor and trochlear palsy suggests a lesion in the cavernous sinus.

A 56-year-old man presented with a complete, pupil-sparing right oculomotor palsy (video E-1). The evaluation was negative, and the palsy resolved within 1 month.

ACKNOWLEDGMENT

The author thanks Dr. Neil Miller for assistance.

REFERENCES

1. Trobe JD. Isolated third nerve palsies. *Sem Neurol* 1986; 6:135–141.
2. Ansons AM, Davis H. Diagnosis and management of ocular motility disorders. 3rd ed. Oxford: Blackwell Science Ltd, 2001:359–360.

Disclosure: The author reports no conflicts of interest.

Neurology®

Teaching Video Is it III alone, or III and IV?

Stephen G. Reich

Neurology 2007;68;E34

DOI 10.1212/01.wnl.0000262060.76472.60

This information is current as of May 21, 2007

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/68/21/E34.full>

Supplementary Material

Supplementary material can be found at:
<http://n.neurology.org/content/suppl/2007/05/20/68.21.E34.DC1>

References

This article cites 1 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/68/21/E34.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All Neuro-ophthalmology

http://n.neurology.org/cgi/collection/all_neuroophthalmology

Diplopia (double vision)

http://n.neurology.org/cgi/collection/diplopia_double_vision

Ocular motility

http://n.neurology.org/cgi/collection/ocular_motility

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.