Obesity and migraine: A population study

To the Editor: We read with interest the recent article by Bigal et al.¹ concerning migraine and obesity. They concluded that obesity is not comorbid to migraine but is significantly associated with the number of headache days, severity, and some clinical features. We would like to introduce our survey of headache in nurses who are working in our hospital.

Almost 700 nurses participated in this survey, and 10% of them have migraine. According to the results, the proportion of migraine subjects with 10 or more headache days per month is nearly 10% in normal or underweight subjects. There was no one whose body mass index was greater than 25 and who had headache of 10 days or more.

We would like the authors to clarify the following points. Are those who are obese in childhood or adolescence at greater risk for developing migraine than those who are not? In addition, overweight people may have sleep apnea syndrome (SAS), hyperglycemia, and hypercholesterolemia. These factors may also play a role in migraine. Migraine is associated with elevated cholesterol level² and mitochondria plays a role in migraine pathogenesis.³

In addition, is it possible that reduction of body weight may improve headache in obese subjects? It would be interesting to investigate whether reduction of body weight in normal or underweight subjects would impact migraine.

Yasuo Iwasaki, Ken Ikeda, *Tokyo, Japan*

Disclosure: The authors report no conflicts of interest.

To the Editor: I read the Bigal et al.¹ article with interest. Is it possible that the increased severity and frequency of migraine demonstrated by the authors in patients with obesity relate to the known relationship between obesity and sleep apnea?

An increased occurrence of migraine headache in patients with obstructive sleep apnea has been demonstrated,^{4,5} so it is possible that the increased frequency of migraine in the authors' obese patients occurred only in those with sleep apnea. The mechanism may operate through hypoxia, and the obese also have an increased susceptibility to hypoxia (e.g., Pickwickian syndrome).

It was recently suggested⁶ that hypoxia may be the common precipitator of migraine, as also reflected in migraine's increased occurrence at altitude⁷ and in patients with arteriovenous malformations (AVM) with reduction in migraine following embolization of the AVM.⁸

Gordon J. Gilbert, *St. Petersburg, FL*

Disclosure: The author reports no conflicts of interest.

Reply from the Authors: We appreciate the opportunity to respond to Drs. Iwasaki, Ikeda, and Gilbert. Although obesity was not a risk factor for migraine in our study, we found that obese migraine sufferers had more frequent, painful and disabling headaches than those who were of normal weight.¹ In separate studies, we also showed that obesity is a risk factor for incident chronic daily headache (headache >15 days per month) and that the association of obesity with transformed migraine is robust while the association of obesity with chronic tension type headache is modest.^{2,9}

Prospective study of symptomatic atherothrombotic intracranial stenoses: The GESICA study

To the Editor: I read with interest the article by Mazighi et al.¹ on the natural history of symptomatic intracranial stenosis. Their population resembles that of the WASID trial regarding the risk factor profile.² However, the reported rates of cerebrovascular events are misleading considering the composite endpoint alone. The annual rate of cerebrovascular events was 19% (38% at 2

Drs. Iwasaki, Ikeda, and Gilbert ask about the potential role of sleep apnea in the exacerbation of migraine in obese subjects. Although we cannot directly test this hypothesis, obesity and snoring are independent risk factors for chronic daily headache.² If snoring is a reasonable proxy for sleep apnea, then the influence of obesity on migraine frequency appears to be independent of sleep apnea. This is an excellent area for further research.

Drs. Iwasaki and Ikeda also identify hyperlipidemia and mitochondrial dysfunction as factors that may confound or contribute to the relationship between migraine and obesity. We do not have the data to test these reasonable hypotheses. Obesity is a proinflammatory state, associated with hyperlipidemia, hypertension, and insulin resistance in the metabolic syndrome.¹⁰ Additional research is required to disentangle the contribution of these features, separately and in combination, to the worsening of migraine and the emergence of chronic daily headache.

Although we recently reviewed mitochondrial dysfunction in migraine,¹¹ we do not know how this is connected with obesity. Nor do we know if the duration of obesity contributes to migraine exacerbations or if weight loss reduces headache frequency or severity.

The identification of factors that exacerbate migraine and contribute to its progression is in its infancy. We look forward to finding more satisfactory answers to the excellent questions raised in these letters.

Marcelo E. Bigal, Richard B. Lipton, *Bronx, NY*

Disclosure: Ortho-McNeil Neurologics funded the analysis of the study discussed in this Correspondence. Drs. Bigal and Lipton have received honoraria from Ortho-McNeil Neurologics unrelated to this study.

Copyright © 2007 by AAN Enterprises, Inc.

References

1. Bigal ME, Liberman JN, Lipton RB. Obesity and migraine: a population study. *Neurology* 2006;66:545–550.
2. Scher AI, Stewart WF, Ricci JA, Lipton RB. Factors associated with the onset and remission of chronic daily headache in a population-based study. *Pain* 2003;106:81–89.
3. Schoenen J. Deficient habituation of evoked cortical potentials in migraine: a link between biology, behavior and trigeminovascular activation? *Biomed Pharmacother* 1996;50:71–78.
4. Loh NK, Dinner DS, Foldvary N, Skobieranda F, Yew WW. Do patients with obstructive sleep apnea wake up with headaches? *Ann Intern Med* 1999;159:1765–1768.
5. Ulfberg J, Carter N, Talback M, Edling C. Headache, snoring, and sleep apnea. *J Neurol* 1996;243:621–625.
6. Gilbert GJ. The purpose of migraine. *Q J Med* 2005;26–27.
7. Silber E, Sonnenberg P, Collier DJ, Pollard AJ, Murdoch DR, Goadsby PJ. Clinical features of headache at altitude. A prospective study. *Neurology* 2003;60:1167–1171.
8. Post MC, Thijs V, Schonewille WJ, et al. Embolization of pulmonary arteriovenous malformations and decrease in prevalence of migraine. *Neurology* 2006;66:202–205.
9. Bigal ME, Lipton RB. Obesity is a risk factor for transformed migraine but not chronic tension-type headache. *Neurology* 2006;67:252–257.
10. Shoelson SE, Lee J, Goldfine AB. Inflammation and insulin resistance. *J Clin Invest* 2006;116:1793–1801.
11. Sparaco M, Feleppa M, Lipton RB, Rapoport AM, Bigal ME. Mitochondrial dysfunction and migraine: evidence and hypotheses. *Cephalalgia* 2006;26:361–372.

years) although the rate of TIA was twice as frequent as stroke (12% vs 7%). Additionally, two thirds of patients who underwent angioplasty did so following a TIA, and the periprocedural rate of neurologic complications was 14%.

The composite endpoint issue is important because one is the apparent magnitude of risk when the composite endpoint is considered alone and the other is the relative importance of the different component endpoints to patients, particularly in a situation

when an invasive procedure becomes a treatment alternative and it is associated with a high periprocedural risk.

When presented with an annual risk of a cerebrovascular event of 19% and a treatment procedure with a complication rate of 14%, one may feel compelled to get the angioplasty. However, if the risk of stroke is considered alone (7%) we might decide for medical treatment. Even when TIA and stroke have the same biological mechanism and the very high early risk of stroke that TIA portends,³ it can be argued that for patients or physicians a TIA may have different importance as outcome.

This problem is illustrated with the VA Cooperative Study on Carotid Endarterectomy.⁴ In that trial, the benefit of the procedure was observed only when the composite endpoint included crescendo TIA and it was outweighed by the risks when TIA was not considered. Some authors recommended that to base a clinical decision on a composite event one has to determine 1) if the component endpoints are of similar importance to patients, 2) if the endpoints occur with similar frequency, and 3) if the expected risk reductions are similar with treatment.

When there are wide variations between the components then the composite endpoint should be abandoned.⁵ Mazighi et al. did not need the addition of TIA to represent the high risk these patients carry. Moreover, death was a better component of a composite endpoint. I commend the authors' restraint on recommending angioplasty or stent placement and acknowledging the need for further clinical trials.

Salvador Cruz-Flores, *St. Louis, MO*

Disclosure: The author reports no conflicts of interest.

Reply from the Authors: We appreciate Dr. Cruz-Flores' comments and agree that the relevance of a composite criterion is certainly inferior to an individual endpoint. In the GESICA study, the recurrence of ischemic events in the territory of a symptomatic intracranial stenosis defined the major endpoint.¹ To what extent TIA and ischemic stroke (IS) are different entities and represent components of a composite endpoint is unclear. They both constitute a recurrent cerebrovascular ischemic event, which is clearly different from a composite endpoint including stroke and myocardial infarction.

TIA and IS share pathogenic mechanisms, and in this specific population of patients with stroke (i.e., symptomatic intracranial stenosis), they are probably part of a continuum. The use of a conventional clinical definition for TIA (which does not include imaging) in the GESICA study suggests that a significant number of TIAs may have been classified as IS if the new definition of TIA is used.⁶ TIAs are often a concern for patients and physicians and they are a serious condition which require adequate therapeutic

decisions. TIAs are an important determinant of stroke, with risks of stroke reported as high as 12% within the first month and looking back after an IS, 23% of patients experience a TIA before their stroke, mainly in the preceding week.^{3,7}

Currently, endovascular treatment remains under investigation. The initial high complications rates are decreasing as a result of interventionists' increasing experience and the development of dedicated devices. The most recent data report periprocedural risks between 5.8% and 9.5%^{8,9} and the long-term follow-up available in patients treated with intracranial angioplasty or stenting shows annual stroke rates ranging between 3.2% and 7.1%.^{1,9} These stroke rates constitute a better outcome compared to the 22% stroke rate in the highest risk cohort in the WASID study.²

However, indications for endovascular intervention in patients with intracranial atherosclerotic remain to be established, but the identification of patients with the highest risk of recurrence of ischemic events will probably maximize the benefit/risk ratio of this technique.

M. Mazighi, X. Ducrocq, S. Bracard, E. Houdart, F. Woimant, *Paris, France*

Disclosure: The authors report no conflicts of interest.

Copyright © 2007 by AAN Enterprises, Inc.

References

1. Mazighi M, Tanasescu R, Ducrocq X, et al. Prospective study of symptomatic atherothrombotic intracranial stenoses: the GESICA study. *Neurology* 2006;66:1187–1191.
2. Chimowitz MI, Lynn MJ, Howlett-Smith H, et al. Comparison of warfarin and aspirin for symptomatic intracranial arterial stenosis. *N Engl J Med* 2005;352:1305–1316.
3. Johnston SC, Gress DR, Browner WS, Sidney S. Short-term prognosis after emergency department diagnosis of TIA. *J Am Med Assoc* 2000;284:2901–2906.
4. Mayberg MR, Wilson E, Yatsu F, et al. Carotid endarterectomy and prevention of cerebral ischemia in symptomatic carotid stenosis. *JAMA* 1991;266:3289–3294.
5. Montori VM, Permyer-Miranda G, Ferreira-González I, et al. Validity of composite end points in clinical trials. *BMJ* 2005;330:594–596.
6. Albers GW, Caplan LR, Easton JD, et al. Transient ischemic attack—proposal for a new definition. *N Engl J Med* 2002;347:1713–1716.
7. Rothwell PM, Warlow CP. Timing of TIAs preceding stroke: time window for prevention is very short. *Neurology* 2005;64:817–820.
8. Henkes H, Miloslavski E, Lowens S, Reinartz J, Liebig T, Kuhne D. Treatment of intracranial atherosclerotic stenoses with balloon dilatation and self-expanding stent deployment (WingSpan). *Neuroradiology* 2005;47:222–228.
9. Marks MP, Wojak JC, Al-Ali F, et al. Angioplasty for symptomatic intracranial stenosis: clinical outcome. *Stroke* 2006;37:1016–1020.

Elderly-onset essential tremor is associated with dementia

To the Editor: We read with great interest the article by Benito-Leon et al.¹ concerning dementia in patients with essential tremor (ET). They found that subjects with onset of ET after age 65 were significantly more likely to be demented than controls, while tremor onset at or below age 65 was not associated with dementia.

While the authors discuss peculiar patterns of Lewy body or AD-type pathology as candidates for the pathologic basis underlying this association, they did not mention the possibility that some of these patients may in fact be carriers of the fragile X premutation (FXP).^{2,3} The fragile X syndrome with mental retardation and typical physical stigmata results from the expansion of a CGG trinucleotide repeat in the FMR1 gene. Normally, FMR1 has 6 to 40 CGG repeats, while the full mutation results from expansions with more than 200 repeats. Those with intermediate expansions are considered as carriers of the FXP which has recently been reported to be associated with the fragile X-associated tremor/ataxia syndrome (FXTAS), including cases misdiagnosed as ET.^{2,4}

FXP carriers frequencies are estimated to be 1 per 260 females and 1 per 760 males with a penetrance to FXTAS close to 50% at the age of 60 and up to 100% for individuals older than 80.³ While cases have been reported in both men and women, FXTAS is

probably more frequent in male subjects who present typically with onset in the sixth decade and including additional executive cognitive deficits, variable degrees of peripheral neuropathy, and occasionally mild parkinsonism.^{2–4}

Although relatively more cases of FXTAS have been found in men and 60.4% of the ET cases in the Benito-Leon et al. study are female, we think that some of these patients may be FXP carriers and FXTAS should be included as an additional alternative possibility to account for this interesting finding.

Renato P. Munhoz, Helio G. Teive, *Curitiba PR Brazil*

Disclosure: The authors report no conflicts of interest.

Reply from the Authors: We thank Drs. Munhoz and Teive for their comments although we do not agree with their idea that FXTAS should be included as an alternative possibility. An occasional patient with FXTAS manifesting as an ET-like picture has been reported by us⁵ and others.^{6,7} These studies included hundreds of ET patients in different cohorts consistently showing that patients with ET do not have a higher than expected rate of FMR1 repeat expansions.⁸ It is unlikely that the increased prevalence of dementia we observed in the patients with ET we studied was due to an increased prevalence of FMR1 repeat expansions in these patients.

We expect that the underlying causes for the dementia in ET will become more apparent with rigorous postmortem studies of ET brains.⁹

Elan D. Louis, MD, MSc, *New York, NY*; Julián Benito-León, MD, PhD, *Madrid, Spain*

The authors report no conflicts of interest.

Copyright © 2007 by AAN Enterprises, Inc.

References

1. Benito-León J, Louis ED, Bermejo-Pareja F, Neurological Disorders in Central Spain Study Group. Elderly-onset essential tremor is associated with dementia. *Neurology* 2006;66:1500–1505.
2. Hagerman PJ, Hagerman RJ. The fragile-X premutation: a maturing perspective. *Am J Hum Genet* 2004;74:805–816.
3. Jacquemont S, Hagerman RJ, Leehey MA, et al. Penetrance of the fragile X-associated tremor/ataxia syndrome in a premutation carrier population. *JAMA* 2004;291:460–469.
4. Leehey MA, Munhoz RP, Lang AE, et al. The fragile X premutation presenting as essential tremor. *Arch Neurol* 2003;60:117–121.
5. García Arocena D, Louis ED, Tassone F, et al. Screen for expanded FMR1 alleles in patients with essential tremor. *Mov Disord* 2004;19:930–933.
6. Tan EK, Zhao Y, Puong KY, et al. Fragile X premutation alleles in SCA, ET, and parkinsonism in an Asian cohort. *Neurology* 2004;63:362–363.
7. Deng H, Jankovic J. Premutation alleles associated with Parkinson's disease and essential tremor. *JAMA* 2004;292:1685–1688.
8. Hall DA, Hagerman RJ, Hagerman PJ, Jacquemont S, Leehey MA. Prevalence of FMR1 repeat expansions in movement disorders. A systematic review. *Neuroepidemiology* 2006;26:151–155.
9. Louis ED, Vonsattel JPG, Honig LS, Ross GW, Lyons KE, Pahwa R. Neuropathological findings in essential tremor. *Neurology* 2006;66:1756–1759.

An almost missed leptomeningeal angioma in Sturge-Weber syndrome

Csaba Juhasz, MD, PhD; and Harry T. Chugani, MD, *Detroit, MI*

Although leptomeningeal angioma is the hallmark of Sturge-Weber syndrome (SWS),¹ some patients with suspected SWS show no MRI evidence of pial contrast enhancement.² A girl with a left port wine stain developed right-sided motor seizures. MRI was performed at 18 months of age. Post-gadolinium T1-weighted images (slice thickness: 2 mm) showed transmedullary enhancement in the left central region (figure) but no clear evidence of leptomeningeal angioma. MRI scanning was continued with a three-dimensional time-of-flight (TOF) MR venography to further evaluate possible venous abnormalities. The axial source images (slice thickness: 0.8 mm) showed left parietal leptomeningeal enhancement (figure) in 20 consecutive planes. Three-dimensional TOF MR venography may increase the chance of detecting a leptomeningeal angioma in children with suspected SWS.

Copyright © 2007 by AAN Enterprises, Inc.

1. Benedikt RA, Brown DC, Walker R, Ghaed VN, Mitchell M, Geyer CA. Sturge-Weber syndrome: cranial MR imaging with Gd-DTPA. *AJNR Am J Neuroradiol* 1993;14:409–415.
2. Fischbein NJ, Barkovich AJ, Wu Y, Berg BO. Sturge-Weber syndrome with no leptomeningeal enhancement on MRI. *Neuroradiology* 1998;40:177–180.

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Csaba Juhasz, Division of Pediatric Neurology/PET Center, Children's Hospital of Michigan, 3901 Beaubien Blvd, Detroit, MI 48201; e-mail: juhasz@pet.wayne.edu

NeuroImages

Figure. Axial MR images taken after gadolinium injection. (A) Early post-gadolinium T1-weighted images showed no evidence of leptomeningeal angioma, only transmedullary veins in the left central region (arrowheads). (B) Post-gadolinium source images from three-dimensional time-of-flight MR venography showed left superior parietal contrast enhancement consistent with a pial angioma (arrows), in addition to visualizing the deep veins.

Neurology[®]

An almost missed leptomeningeal angioma in Sturge-Weber syndrome

Csaba Juhasz and Harry T. Chugani

Neurology 2007;68;243

DOI 10.1212/01.wnl.0000242581.43024.0a

This information is current as of January 15, 2007

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/68/3/243.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2007/07/27/68.3.243.DC1
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/68/3/243.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): MRI http://n.neurology.org/cgi/collection/mri Other neurocutaneous disorders http://n.neurology.org/cgi/collection/other_neurocutaneous_disorders
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.