Neuroimaging of retinal nerve fiber layer in AD using optical coherence tomography

Figure Normal retinal NFL (A) and moderate AD NFL (B)

On the left are optical coherence tomography images; to the right are graphs depicting the thickness. The arrows point to the superior area of the NFL and indicate areas of difference due to AD. The normal has a superior NFL thickness of 104 μm and the AD thickness is 58 μm . AD = Alzheimer disease; NFL = nerve fiber layer; TEMP = temporal; SUP = superior; NAS = nasal; INF = inferior. Images were acquired using a Stratus OCT™ (Carl Zeiss Meditec, Inc.).

Optical coherence tomography (OCT) is used for the management of diseases affecting the eye. It has the ability to measure the thickness of the nerve fiber layer (NFL) of the retina. OCT is similar to ultrasound, but utilizes light. Unlike ultrasound, there is no tissue contact. OCT measures structures less than 10 μm scale. In a study that evaluated individuals with Alzheimer disease (AD) and control individuals, Danesh-Meyer and colleagues¹ found a reduction in NFL in those with AD, but not control individuals. Parisi and colleagues² also found significant reductions in the NFL of those with AD and this group included mild AD.

Denise A. Valenti, OD, FAAO, Boston, MA

Address correspondence and reprint requests to Dr. D.A. Valenti, Ophthalmology, Boston University, 715 Albany St., 9th fl., Boston, MA 02118; dvalenti@bu.edu

Supported by NIH Disability Supplement to R01 1AG15361, Mentored Research/Training: Changes in Optic Nerve Fiber Layer Secondary to Alzheimer's Disease.

Disclosure: The author reports no conflicts of interest.

The author thanks Thomas Laudate for his assistance with the acquisition of images in the Vision and Cognition Laboratory at Boston University.

REFERENCES

1. Danesh-Meyer HV, Birch H, Ku JY, Carroll S, Gamble G. Reduction of optic nerve fibers in patients with Alzheimer disease identified by laser imaging. *Neurology* 2006;67:1852–1854.
2. Parisi V, Restuccia R, Fattapposta F, Mina C, Bucci MG, Pierelli F. Morphological and functional retinal impairment in Alzheimer's disease patients. *Clin Neurophysiol* 2001;112:1860–1867.

Neurology®

Neuroimaging of retinal nerve fiber layer in AD using optical coherence tomography

Denise A. Valenti

Neurology 2007;69;1060

DOI 10.1212/01.wnl.0000280584.64363.83

This information is current as of September 4, 2007

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/69/10/1060.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/69/10/1060.full#ref-list-1>

Citations

This article has been cited by 2 HighWire-hosted articles:
<http://n.neurology.org/content/69/10/1060.full##otherarticles>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.