Nitromethane encephalopathy MRI

Figure T2-weighted fluid attenuated inversion recovery axial (A, B), coronal (C), and sagittal (D) sections showing bilateral and symmetric hyperintense lesions in cerebellar white matter, tonsils, uvula, and tectum

These lesions were less conspicuous on T2-weighted images. Symmetric lesions support the diagnosis of toxic encephalopathy.

After nitromethane fuel ingestion, an 18-year-old male mechanic presented with generalized tonic-clonic seizures progressing to partial motor status epilepticus. At physical examination hypertension was noted. Symptomatic therapy was initiated, including valproic acid, β -blockers, and clonidine. When neurologic examination could be performed, it showed mild left dysmetria in finger-to-nose testing, low-frequency intention tremor, broad-based gait, and inability to tandem-walk. MRI (figure) demonstrated bilateral and symmetric lesions in cerebellar white matter, tonsils, uvula, and colliculi. Valproic acid was maintained. Clinical and MRI 8-month follow-up showed no abnormalities. To our knowledge, this is the first documented case of reversible encephalopathy by nitromethane ingestion.^{1,2}

Elena Alventosa Fernández, MD, Candelaria González González, MD, Javier Crisóstomo Pardillo, MD, Vicente Martín García, MD, Santa Cruz de Tenerife, Spain

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Elena Alventosa Fernández, C/Santa Rosalía, 5-3°, 38002 Santa Cruz de Tenerife, Spain; elena.alventosa@gmail.com814814

1. Sclar G. Encephalomyeloradiculoneuropathy following exposure to an industrial solvent. *Clin Neurol Neurosurg* 1999; 101:199–202.
2. Booth C, Naidoo D, Rosengerg A, Kainer G. Elevated creatinine after ingestion of model aviation fuel: interference with the Jaffe reaction by nitromethane. *J Paediatr Child Health* 1990;35:503–504.

Neurology[®]

Nitromethane encephalopathy MRI

Elena Alventosa Fernández, Candelaria González González, Javier Crisóstomo Pardillo, et al.

Neurology 2008;70;814

DOI 10.1212/01.wnl.0000304254.18355.b3

This information is current as of March 3, 2008

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/70/10/814.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/70/10/814.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/70/10/814.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): MRI http://n.neurology.org/cgi/collection/mri Other toxicology http://n.neurology.org/cgi/collection/other_toxicology Solvents http://n.neurology.org/cgi/collection/solvents
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.