Understanding drug-induced parkinsonism

Separating pearls from oy-sters

Maria Victoria G.
Alvarez, MD
Virgilio Gerald H.
Evidente, MD

Address correspondence and
reprint requests to Dr. Maria
V.G. Alvarez, Mayo Clinic
Arizona, 13400 Shea Blvd.,
Scottsdale, AZ 85259
martyzqt@aol.com

In 1817, James Parkinson described the shaking palsy now known as Parkinson disease (PD). Descriptions evolved until the term “parkinsonism” now refers to a syndrome characterized by the presence of tremor, rigidity, and bradykinesia in addition to loss of postural reflexes and freezing. The most common cause of parkinsonism is PD. However, in any parkinsonian patient, one must obtain a careful medical and medication history, as drug-induced parkinsonism (DIP) is often reversible, especially if the offending drug is discontinued early.

CASE SCENARIO A 46-year-old man developed upper limb tremors 2 weeks after initiating perphenazine for mood disorder. Perphenazine was promptly changed to quetiapine, though his tremor persisted. He had a history of hypertension, hypercholesterolemia, and chronic nausea. His medications include verapamil, lovastatin, and metoclopramide.

Motor, sensory, and cerebellar testing were normal. Although mildly stooped, he had normal gait and postural reflexes. Extrapyramidal examination revealed a mild chin tremor, moderate symmetric 3 to 4 Hz resting tremor of the upper and lower limbs, and faster frequency postural tremor of both arms. Finger tapping was moderately irregular bilaterally.

“PEARLS” FOR EVALUATING DIP Many patients with DIP are often misdiagnosed with PD. While recognizing clues that suggest DIP is critical (see table 1A),¹ identifying risk factors (table 1B)² and offending agents is just as important (summarized in table 2).³ DIP falls under the category of secondary parkinsonism (due to drugs, toxins, structural lesions).⁴ Thus, early recognition may lead to a better prognosis.

The most common culprits are the neuroleptics with potent dopamine D2-receptor blocking actions, including haloperidol and perphenazine.¹ If chronic neuroleptic drug use is required, the lowest effective dose or an atypical neuroleptic drug is preferred. In one study, clozapine was shown to significantly improve psychosis without worsening or causing motor symptoms of parkinsonism.⁵ However, clozapine can cause agranulo-

Table 1 Pearls regarding drug-induced parkinsonism (DIP)

Clues that might suggest DIP
Subacute bilateral onset and progression of symptoms time-locked with medication intake
Early presence of postural tremor
Concurrent oral-buccal dyskinesias
Risk factors for development of DIP
High dose, high potency, piperazine side chain neuroleptics
Elderly
Female (female:male ratio 2:1)
Hereditary Parkinson disease
Preclinical parkinsonism
AIDS
Coexistence of tardive dyskinesia

cytosis, which requires regular monitoring of blood count.

Metoclopramide is used widely in patients with gastric motility disorders and it has a chemical structure similar to the neuroleptic drug chlorpromazine.¹ The occurrence of parkinsonism secondary to chronic metoclopramide use had been well documented.¹ Another neuroleptic, prochlorperazine, is frequently used to treat gastric motility disorder. It too has the potential to cause parkinsonism and can cause acute dystonia in the first 24 to 48 hours.¹

SKIP THE “OY-STERS”: AVOIDING PITFALLS

Nonparkinsonian tremors. It is important to tell the patient that tremor does not automatically mean PD. In fact, there are several types of tremor associated with different diseases and disorders. Essential tremor (ET) has a gradual onset of 4 to 12 Hz postural upper limb tremor (less commonly head or voice), with no other neurologic signs except for cogwheeling.⁶ Enhanced physiologic tremor (EPT) is a postural, fast frequency upper limb tremor. EPT is the most common drug-induced tremor and many

From Mayo Clinic Arizona, Scottsdale.

Disclosure: The authors report no conflicts of interest.

Copyright © 2008 by AAN Enterprises, Inc.

Table 2 Potential culprits or causes of drug-induced parkinsonism (DIP)

High risk	
Dopamine D2-receptor blockers	Neuroleptics: butyrophenones ³ (haloperidol and others), phenothiazines ³ (prochlorperazine), thioxanthenes ³ (thiothixene), dibenzoxazine ³ (loxapine), others
	Atypical neuroleptics: risperidone ³ (especially in higher concentration)
	Antiemetics/gastric motility agents: substituted benzamides ^{1,3} (metoclopramide, prochlorperazine, and others)
Dopamine depleters	Tetrabenazine ³
Antihypertensive agents thought associated with DIP by reducing dopamine levels	Reserpine and alpha-methyldopa ^{1,3}
Intermediate risk	
Calcium channel blockers with dopamine antagonist activity	Flunarizine, ^{1,3} cinnarizine, ³ verapamil ¹
Certain anticonvulsants	Valproate ^{3*}
Mood stabilizer	Lithium ³ (causes tremor and myoclonus)
Atypical neuroleptics	Risperidone, ³ clozapine, ⁵ and others (especially in higher dose)
Lower risk [†]	
Antihypertensives	Diltiazem, ³ captopril ³
Antiarrhythmic	Amiodarone, ^{1,3} procaine ³
Immunosuppressants	Cyclosporine, ^{1,3} tacrolimus ³
Antidepressants	Fluoxetine ³ (and other SSRIs), tricyclic antidepressants, ³ and certain MAOIs, e.g., phenelzine ³
Antifungals	Co-trimoxazole, amphotericin B ³
Antibiotics	Trimethoprim-sulfamethoxazole ³
Antivirals	Vidarabine, acyclovir (and antiretroviral drugs for HIV) ³
Chemotherapeutics	Thalidomide, cytarabine, ifosfamide, vincristine, tamoxifen, and cytosine arabinoside ^{1,3}
Statins	Lovastatin and others ^{1,3}
Hormones	Levothyroxine, ³ medroxyprogesterone, ³ epinephrine ³
Others	Bethanechol, ³ pyridostigmine, ³ donepezil ³

*There are anecdotal reports of tremor associated with some newer anticonvulsants, though parkinsonism was not described. Examples are tiagabine, gabapentin, oxcarbazepine monotherapy, and lamotrigine.³

[†]Medications that rarely can cause or worsen parkinsonism and tremor.

MAOI = monoamine oxidase inhibitor; SSRI = selective serotonin reuptake inhibitor.

patients have previously unnoticed, undiagnosed tremors.³ EPT can also be associated with endocrine dysfunction, particularly hyperthyroidism, hypoglycemia, and caffeine intake.³ Cerebellar tremor is a slow intention tremor (usually less than 5 Hz) and may be postural.⁷

Tests you do not want to miss. Apart from identifying a potential offending agent, it is also a must to look for other secondary causes of parkinsonism. Wilson disease (WD) should be considered in patients with onset of movement disorder before age 50 years. It is critical not to miss the diagnosis of WD because it is usually treatable with drugs that chelate copper.³ Clinical manifestations may include hepatic, neurologic, psychiatric, and oph-

thalmologic abnormalities.⁸ The classic neurologic sign is a wing-beating proximal upper limb postural tremor, though parkinsonism can be seen. Screening tests include serum ceruloplasmin (normal >20 mg/dL), 24-hour urine copper (normal <100 pg/24 hours), and slit-lamp eye examination looking for copper deposits within the cornea around the iris (Kayser-Fleischer rings).³

Hypothyroidism can present with subacute to chronic onset of symmetric parkinsonism. Serum TSH may be a useful screening test.

MANAGING DIP DIP may persist or remit slowly despite prompt discontinuation of the offending drug. Some patients may require medications temporarily to relieve symptoms.

Pearls in choosing drugs. Use of L-dopa or anticholinergic agents may be indicated and effective.¹ Symptoms should eventually resolve if the parkinsonism was drug-induced.¹ In our experience, levodopa and dopamine agonists can potentially improve most features of parkinsonism, though they may be less effective in alleviating severe neuroleptic-induced tremor.

Avoiding the oysters. Tetrabenazine has been used worldwide to alleviate hyperkinetic movements such as tardive dystonia and chorea, but it can worsen parkinsonism.³ Perhaps this can be explained best by its dopamine-depleting property. Beta-blockers like metoprolol have been regarded as mainstays of pharmacologic therapy for ET, but not for DIP.⁹ Similarly, primidone has been useful in reducing tremor in ET but not in DIP.⁹ Surgical procedures are available that effectively ameliorate tremor in ET that is refractory to medical management.⁹ Their potential effectiveness in DIP remains to be investigated.

Though our patient discontinued perphenazine, his persistent tremor probably was due to continued intake of metoclopramide. He experienced 50% improvement after discontinuing metoclopramide and being placed on 750 mg/day of levodopa.

PEARLS TO TAKE HOME Prompt recognition of DIP is the key. The next step is discontinuation of the culprit. Most cases remit within 4 months after stopping the causative drugs.¹ Rarely, parkinsonism can be persistent, unremitting, and disabling. In such cases, certain drugs may provide symptomatic relief. Finally, prevention remains the most important strategy.²

REFERENCES

- Adler CH. Differential diagnosis of Parkinson's disease. *Med Clin N Am* 1999;83:349-367.

2. Sethi KD. Movement disorders induced by dopamine blocking agents. *Semin Neurol* 2001;21:59–68.
3. Morgan JC, Sethi KD. Drug-induced tremors. *Lancet Neurol* 2005;4:866–876.
4. Ahlskog JE. Diagnosis and differential diagnosis of Parkinson's disease and parkinsonism. *Parkinsonism Relat Disord* 2000;7:63–70.
5. The Parkinson Study Group. Low-dose clozapine for the treatment of drug-induced psychosis in Parkinson's disease. *N Engl J Med* 1999;340:757–763.
6. Singer C, Sanchez-Ramos J, Weiner WJ. Gait abnormality in essential tremor. *Mov Disord* 1994;9:193–196.
7. Mark MH. Tremor disorders. *Continuum* 2005;10:113–127.
8. Pfeiffer RF. Wilson's disease. *Semin Neurol* 2007;27:123–132.
9. Sullivan KL, Hauser RA, Zesiewicz TA. Essential tremor: epidemiology, diagnosis, and treatment. *Neurologist* 2004;10:250–258.

Neurology[®]

Understanding drug-induced parkinsonism: Separating pearls from oysters

Maria Victoria G. Alvarez and Virgilio Gerald H. Evidente

Neurology 2008;70:e32-e34

DOI 10.1212/01.wnl.0000302255.49113.51

This information is current as of May 21, 2008

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/70/8/e32.full>

References

This article cites 9 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/70/8/e32.full#ref-list-1>

Citations

This article has been cited by 3 HighWire-hosted articles:
<http://n.neurology.org/content/70/8/e32.full##otherarticles>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.