Subarachnoid fat

Unusual migration from pelvis to brain

Figure Subarachnoid migration of fatty material from a pelvic fracture

Fat droplets (white arrows) appearing hypodense on transverse CT (A) and hyperintense on sagittal T1-weighted MRI (B) were demonstrated within cerebral ventricles and subarachnoid spaces. Coronal oblique T1-weighted (C) and STIR (D) MRI as well as transverse CT (E, F) revealed a sacral fracture (arrowheads) extending through an S2 Tarlov cyst (asterisk) containing a small amount of fat (black arrow).

A 77-year-old man was admitted after a fall in which he struck his head and pelvis. Neurologic examination was unremarkable. CT and MRI scans of the head (figure, A and B) revealed fat droplets in the cerebral ventricles and subarachnoid spaces. CT and MRI scans of the pelvis (figure, C–F) demonstrated a sacral fracture extending into a Tarlov (perineurial) cyst within the S2 foramen.

We hypothesize that fatty bone marrow migrated from the fracture site to brain through a meningeal breach at the Tarlov cyst. No specific treatment was required.

Dissemination of free fatty material from the spinal subarachnoid space to the brain has been previously reported in a patient with spina bifida occulta following a traumatic sacral pseudomeningocele¹ and following the rupture of spinal dermoid tumors.²

T. Moser, MD, D. Szwarc, MD, G. Zöllner, MD, S. Vinzio, MD, J.-L. Dietemann, MD, and S. Kremer, MD, PhD, Strasbourg, France

Disclosure: The authors report no disclosures.

Address correspondence and reprint requests to Dr. T. Moser, Department of Radiology, Hôpital de Hautepierre, CHU Strasbourg, 67098, France; thomas.moser@chru-strasbourg.fr

1. Uff C, Bradford R. Retrograde intraventricular hemorrhage caused by a traumatic sacral pseudomeningocele in the presence of spina bifida occulta: case report. *J Neurosurg Spine* 2005;3:390–392.
2. Calabro F, Capellini C, Jinkins JR. Rupture of spinal dermoid tumors with spread of fatty droplets in the cerebrospinal fluid pathways. *Neuroradiology* 2000;42:572–579.

Neurology®

Subarachnoid fat: Unusual migration from pelvis to brain

T. Moser, D. Szwarc, G. Zöllner, et al.

Neurology 2008;71;1838

DOI 10.1212/01.wnl.0000335939.92361.b1

This information is current as of November 24, 2008

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/71/22/1838.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/71/22/1838.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Cerebrospinal Fluid

http://n.neurology.org/cgi/collection/cerebrospinal_fluid

CT

<http://n.neurology.org/cgi/collection/ct>

MRI

<http://n.neurology.org/cgi/collection/mri>

Other trauma

http://n.neurology.org/cgi/collection/other_trauma

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.