Interictal hypermetabolic subcortical band on brain FDG-PET in *doublecortin* mutation

Figure 1 Brain MRI revealing subcortical band heterotopia

T2-weighted brain MRI (A) shows subcortical band heterotopia. Compared to basal ganglion on a 10% point color scale, FDG-PET (B) shows intensely increased FDG uptake (arrows) by the subcortical band and reduced uptake by the cerebral cortex (arrowheads).

Figure 2 Interictal scalp EEG with generalized slow spike and wave (100–200 mV, 1.5–2.5 Hz, in runs of 2–5 seconds) maximum in the parieto-occipital (80%, illustrated in figure) and frontal (20%) regions present during 70% of record

A 13-year-old girl with generalized tonic and grand mal seizures had brain MRI that revealed subcortical band heterotopia (SBH) (figure 1). Twenty-four-hour EEG showed pervasive generalized slow spike and wave pattern (figure 2). Blood DNA confirmed a mutation in the *doublecortin* (*DCX*) gene. Interictal 18-fluoro-deoxy (FDG) PET showed reduced FDG uptake (~10%) in the cerebral cortex and intensely increased uptake (~20%) in the subcortical band (figure 1), suggesting epileptogenic subcortical band correlating with pervasive slow spike wave pattern on simultaneous EEG. Two PET reports^{1,2} showed similar or minor increased FDG uptake in the heterotopic band compared to the cerebral cortex.

Joanna Fong, MD, Guiyun Wu, MD, Elaine Wyllie, MD, Ajay Gupta, MD, Cleveland, OH

Disclosure: The authors report no disclosures.

Address correspondence and reprint requests to Dr. Ajay Gupta, S-51, Epilepsy Center/Neurological Institute, Cleveland Clinic Foundation, Cleveland, OH 44195; guptaaj1@ccf.org

1. Miura K, Watanabe K, Maeda N, et al. Magnetic resonance imaging and positron emission tomography of band heterotopia. *Brain Dev* 1993;15:288–290.
2. De Volder AG, Gadisseux JF, Michel CJ, et al. Brain glucose utilization in band heterotopia: synaptic activity of “double cortex”. *Pediatr Neurol* 1994;11:290–294.

Neurology[®]

Interictal hypermetabolic subcortical band on brain FDG-PET in *doublecortin* mutation

Joanna Fong, Guiyun Wu, Elaine Wyllie, et al.

Neurology 2008;71;535

DOI 10.1212/01.wnl.0000325014.43936.6a

This information is current as of August 11, 2008

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/71/7/535.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/71/7/535.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All Epilepsy/Seizures

http://n.neurology.org/cgi/collection/all_epilepsy_seizures

EEG

http://n.neurology.org/cgi/collection/eeg_

EEG; see Epilepsy/Seizures

http://n.neurology.org/cgi/collection/eeg_see_epilepsy-seizures

PET

<http://n.neurology.org/cgi/collection/pet>

PET in epilepsy

http://n.neurology.org/cgi/collection/pet_in_epilepsy

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.