Teaching NeuroImages:

Sinus pericranii

Antonetta Margaretha
Gezina Sas, MD
Fop van Kooten, PhD

Address correspondence and
reprint requests to Dr. F. van
Kooten, Department of
Neurology, Erasmus Medical
Centre, PO Box 2040, 3000 CA
Rotterdam, The Netherlands
f.vankooten@erasmusmc.nl

An 18-year-old woman was referred for a slowly growing tumor on her scalp, which had been present since her birth. Examination revealed a soft-tissue non-pulsating mass on her scalp (figure 1) which became larger in recumbent position and with Valsalva maneuver. CT showed that almost the entire venous system drained through enlarged parietal foramina with dilated emissary veins connected to multiple subcutaneous veins, which is called sinus pericranii (SP) (figure 2). Generally, SP is located frontally in

Figure 1 Skull x-ray showing the tumor (T) in the parieto-occipital region of the skull

Figure 2 CT scan without (A and B) and with contrast (C and D) just beneath (A and C) and just above (B and D) the skull

CT scan without (A and B) and with contrast (C and D) with axial slices intracranially just under (A and C) and extracranially just above (B and D) the skull through the tumor, showing bilateral enlarged parietal foramina (F) with dilated emissary veins and the varicosity of the tumor. Not shown here, but visible on CT, were hypoplasia of the falx, bilateral small transverse and sigmoid sinuses, and small jugular foramina with minimally developed jugular veins.

the midline and connected with the superior sagittal sinus. SP is commonly asymptomatic, and classified as dominant if the major venous flow is through the SP and accessory if it concerns a minor part of the venous flow. The prognosis is nearly always good with a low risk of bleeding. Only accessory SPs can be treated safely by surgical intervention or endovascular embolization.^{1,2} Treatment is not recommended for dominant SP and we did not recommend treatment in this case.

REFERENCES

1. Gandolfo C, Krings T, Alvarez H, et al. Sinus pericranii: diagnostic and therapeutic considerations in 15 patients. *Neuroradiology* 2007;49:505–514.
2. Kaido T, Kim YK, Ueda K. Diagnostic and therapeutic considerations for sinus pericranii. *J Clin Neurosci* 2006; 13:788–792.

From the Department of Neurology, Erasmus Medical Centre, Rotterdam, The Netherlands.

Disclosure: The authors report no disclosures.

Neurology®

Teaching NeuroImages: Sinus pericranii

Antonetta Margaretha Gezina Sas and Fop van Kooten

Neurology 2009;72:e66

DOI 10.1212/01.wnl.0000345672.15367.dc

This information is current as of April 6, 2009

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/72/14/e66.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/72/14/e66.full#ref-list-1>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://n.neurology.org/content/72/14/e66.full##otherarticles>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
All Clinical Neurology
http://n.neurology.org/cgi/collection/all_clinical_neurology
Arteriovenous malformation
http://n.neurology.org/cgi/collection/arteriovenous_malformation
CT
<http://n.neurology.org/cgi/collection/ct>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.