Teaching Video NeuroImages: Dancing epilepsy

Ritu Bagla, MD
John S. Khoury, MD
Christopher Skidmore,
MD

Address correspondence and
reprint requests to Dr. John
Khoury, 900 Walnut St., Suite
200, Philadelphia, PA 19107
Jskhoury@gmail.com

Figure 1 Fluid-attenuated inversion recovery coronal MRI revealed encephalomalacia and gliosis in the left frontal and right anterior temporal regions

A 39-year-old, right-handed man with refractory complex partial seizures after head trauma at age 15 (figure 1) was admitted for video-EEG monitoring. His girlfriend describes dancing movements with unresponsiveness lasting a few minutes (see video). The interictal EEG was suggestive of a left temporal focus (figure 2); however, the ictal EEG was poorly localized and the overall localization was not determined. The dancing semiology is not typical of temporal lobe epilepsy, since automatisms involving proximal muscles and complex behaviors, such as bicycling, are suggestive of frontal lobe epilepsy.¹ Dancing is another behavioral manifestation of epilepsy that has not been previously reported.

REFERENCE

1. Kotagal P, Arunkumar G, Hammel J, Mascha ED. Complex partial seizures of frontal lobe onset: statistical analysis of ictal semiology. *Seizure* 2003;12:268–281.

Figure 2 Interictal EEG

Interictal left anterior temporal sharp waves. Sensitivity 150 μ V/mm.

Supplemental data at
www.neurology.org

From Thomas Jefferson University Hospital (R.B., J.S.K., C.S.) and Thomas Jefferson Medical College (C.S.), Philadelphia, PA.

Disclosure: Dr. Skidmore has the following conflicts of interest to disclose for research support: UCB Pharma, Ortho McNeil, Schwarz, Medtronic, Neuropace, Ovation, Marinus, NIH. He is a consultant for the Amarin Corporation and a speaker for UCB Pharma and Ortho McNeil.