Ageotropic central positional nystagmus in nodular infarction

Figure Diffusion-weighted axial (A) and sagittal (B) and T2-weighted coronal (C) MRI disclose isolated nodular infarction (arrow). The inset indicates the imaging plane of the corresponding image

A 36-year-old man with a history of hypertension and atrial fibrillation presented with acute vertigo and imbalance. Neurologic examination revealed ataxic gait and ageotropic positional nystagmus during head-turning while supine (video on the *Neurology*[®] Web site at www.neurology.org). Other findings of the neurologic examination were normal. MRI showed bilateral nodular infarction (figure).

Central positional nystagmus mostly develops in lesions involving the brainstem or cerebellum.¹ Isolated nodular infarction may show spontaneous horizontal, periodic alternating, post-head-shaking, and positional nystagmus in addition to imbalance.² Ageotropic central positional nystagmus may be a manifestation of nodular infarction and should be differentiated from benign positional vertigo involving the horizontal semicircular canal.²

Jungmoo Nam, MD, Seonhye Kim, MD, Youngeun Huh, MD, and Ji Soo Kim, MD, PhD, Seoul, Republic of Korea

Supported by a grant of the Korea Health 21 R&D Project, Ministry of Health & Welfare, Republic of Korea (A080750).

Disclosure: Dr. Nam, Dr. S. Kim, and Dr. Huh report no disclosures. Dr. J.S. Kim serves as an Associate Editor of the *Journal of Korean Neurological Association* and on the editorial board of the *Journal of Korean Society for Clinical Neurophysiology*.

Address correspondence and reprint requests to Dr. Ji Soo Kim, Department of Neurology, Seoul National University College of Medicine, Seoul National University Bundang Hospital, 300 Gumi-dong, Bundang-gu, Seongnam-si, Gyeonggi-do, 463-707, Korea; jisookim@snu.ac.kr

Supplemental data at
www.neurology.org

1. Büttner U, Helmchen C, Brandt T. Diagnostic criteria for central versus peripheral positioning nystagmus and vertigo: a review. *Acta Otolaryngol* 1999;119:1–5.
2. Moon IS, Kim JS, Choi KD, et al. Isolated nodular infarction. *Stroke* 2009;40:487–491.

Neurology®

Ageotropic central positional nystagmus in nodular infarction

Jungmoo Nam, Seonhye Kim, Youngeun Huh, et al.

Neurology 2009;73;1163

DOI 10.1212/WNL.0b013e3181bacfde

This information is current as of October 5, 2009

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/73/14/1163.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2009/10/04/73.14.1163.DC1
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/73/14/1163.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/73/14/1163.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Infarction http://n.neurology.org/cgi/collection/infarction Nystagmus http://n.neurology.org/cgi/collection/nystagmus Vertigo http://n.neurology.org/cgi/collection/vertigo
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.