Hemorrhagic Wernicke encephalopathy in a patient with liver transplantation

Figure A 40-year-old woman with Wernicke encephalopathy

Initial CT (A), fluid-attenuated inversion recovery (FLAIR) (B), and T1-weighted MRI (C) at the level of midbrain show early subacute stage hemorrhage in bilateral colliculi. FLAIR at the level of basal ganglia (D) shows symmetric signal increase in medial thalami (arrows). Note hydrocephalus in all initial images. Follow-up CT after 20 days (E) shows resolving hemorrhage in colliculi. Fifteen days later, MRI (F-H) shows resolution of hemorrhage, hemosiderin deposition in colliculi, and reversibility of thalamic signal change and hydrocephalus.

A 40-year-old woman who received a liver transplantation developed nystagmus, downward gaze palsy of her right eye, and gradual mental deterioration. CT and MRI (figure) showed signal change in medial thalami and hemorrhage in the bilateral inferior colliculi, suggesting Wernicke encephalopathy. Swelling around the aqueduct resulted in hydrocephalus. After beginning thiamine supplementation (100 mg/day), her consciousness began to return. Follow-up MRI revealed partially resolved hemorrhages and improved hydrocephalus. Gross hemorrhage is an uncommon finding in Wernicke encephalopathy, although microscopic hemorrhage is well-documented in pathology textbooks.¹ Typical findings include signal change in mamillary bodies, medial thalami, tectal plate, and periaqueductal gray matter.²

Na Young Shin, MD, Hyo Suk Nam, MD, and Seung-Koo Lee, MD, Seoul, Korea

Disclosure: The authors report no disclosures.

Address correspondence and reprint requests to Dr. Seung-Koo Lee, Department of Radiology, Yonsei University College of Medicine, 250 Sungsanro, Seodaemun-gu, Seoul 120-752, Korea; slee@yuhs.ac

1. Vortmeyer AO, Hagel C, Laas R. Haemorrhagic thiamine deficient encephalopathy following prolonged parenteral nutrition. *J Neurol Neurosurg Psychiatry* 1992;55:826–829.
2. Roh JH, Kim JH, Koo Y, et al. Teaching *NeuroImage*: Diverse MRI signal intensities with Wernicke encephalopathy. *Neurology* 2008;70:e48.

Neurology®

Hemorrhagic Wernicke encephalopathy in a patient with liver transplantation

Na Young Shin, Hyo Suk Nam and Seung-Koo Lee

Neurology 2009;73;1423

DOI 10.1212/WNL.0b013e3181bd82b9

This information is current as of October 26, 2009

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/73/17/1423.full>

References

This article cites 2 articles, 2 of which you can access for free at:
<http://n.neurology.org/content/73/17/1423.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

CT

<http://n.neurology.org/cgi/collection/ct>

MRI

<http://n.neurology.org/cgi/collection/mri>

Nutritional

<http://n.neurology.org/cgi/collection/nutritional>

Nystagmus

<http://n.neurology.org/cgi/collection/nystagmus>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.