Diffuse hyperintense brainstem lesions in neuroborreliosis

Figure 1 Brain MRI at admission: Axial fluid-attenuated inversion recovery images show extensive diffuse hyperintense lesions with mild swelling in the diencephalon and mesencephalon (A, B) and focal cerebellar lesions (A)

Figure 2 Brain MRI at 6 months' follow-up: Almost complete resolution of abnormalities after 6 months (A, B), but certain low-density areas are noticeable (A)

A 64-year-old woman presented with a 2-month history of neck pain, wasting, and fatigue followed by gait disturbance, dysarthria, and dysmetria. MRI showed profound hyperintensities in diencephalon, mesencephalon, and cerebellum correlating with symptoms (figure 1). Lymphoma, vasculitis, multiple sclerosis, and tuberculosis were excluded. Neuroborreliosis was diagnosed based on intrathecal immunoglobulin G synthesis (specific antibody index¹ by indirect immunofluorescence 6.9, normal <2) along with abnormal CSF (113 cells/mm³, protein 2.25 g/L, positive oligoclonal bands).

The specific antibody index is calculated as follows:

$$\frac{\text{titer of IgG in CSF}}{\text{titer of IgG in serum}} : \frac{\text{total concentration of IgG in CSF}}{\text{total concentration of IgG in serum}} = \frac{1:1024}{1:2048} : \frac{636 \text{ mg/L}}{8800 \text{ mg/L}}$$

Intravenous ceftriaxone led to nearly complete resolution of signs, symptoms, and MRI abnormalities (figure 2), demonstrating an exceptional case in contrast to usually unspecific MRI findings in neuroborreliosis.²

Adrian Haene, MD, Mathias Tröger, MD, Aarau, Switzerland

Disclosure: The authors report no disclosures.

Address correspondence and reprint requests to Dr. Adrian Haene, Department of Neurology, Cantonal Hospital Aarau, Tellstrasse, 5001 Aarau, Switzerland; adrian.haene@swissonline.ch

1. Blanc F, Jaulhac B, Fleury M, et al. Relevance of the antibody index to diagnose Lyme neuroborreliosis among seropositive patients. *Neurology* 2007;69:953–958.
2. Aalto A, Sjöwall J, Davidsson L, Forsberg P, Smedby Ö. Brain magnetic resonance imaging does not contribute to the diagnosis of chronic neuroborreliosis. *Acta Radiol* 2007;48:755–762.

Neurology®

Diffuse hyperintense brainstem lesions in neuroborreliosis

Adrian Haene and Mathias Tröger

Neurology 2009;73;326

DOI 10.1212/WNL.0b013e3181af78cc

This information is current as of July 27, 2009

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/73/4/326.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/73/4/326.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.