Sleep-specific atrioventricular block

Figure Sleep study: Asleep (A) and awake (B)

A 39-year-old man presented to the sleep laboratory for evaluation of intermittent difficulty sleeping. His sleep study was essentially normal, with exception of the electrocardiographic tracing (ECG), which demonstrated second-degree atrioventricular block (figure, A, and detail in C). A screening 12-lead ECG was normal, when obtained when the patient was awake. The patient had no cardiac symptoms.

Strikingly, this arrhythmia was present continuously, regardless of sleep stage, except during the brief times awake in that night's study (figure, B). Therefore, this finding is likely sleep dependent, rather than resulting from circadian influences. (For a review of the interaction between circadian biology and a variety of cardiac arrhythmias, see reference 1.) In fact, one could reliably discern whether the patient was awake or asleep throughout the night, simply by looking for the presence or absence of conduction block on the ECG. This finding widens the scope of arrhythmias seen in sleep, currently reported in severe apnea,² and in REM sleep.³

Jeffrey M. Ellenbogen, MMSc, MD, Boston, MA

Disclosure: Dr. Ellenbogen has received travel expenses and/or honoraria for lectures or educational activities not funded by industry.

Address correspondence and reprint requests to Dr. Jeffrey M. Ellenbogen, Division of Sleep Medicine, Neurology Department, Mass General Hospital, 55 Fruit Street, Wang ACC 720, Boston, MA 02114; jeffrey_ellenbogen@hms.harvard.edu

1. Portaluppi F, Hermida RC. Circadian rhythms in cardiac arrhythmias and opportunities for their chronotherapy. *Adv Drug Deliv Rev* 2007;59:940–951.
2. Harbison J, O'Reilly P, McNicholas WT. Cardiac rhythm disturbances in the obstructive sleep apnea syndrome: effects of nasal continuous positive airway pressure therapy. *Chest* 2000;118:591–595.
3. Kawana F, Kasai T, Maeno K, Momomura S, Narui K. Atrioventricular block during the phasic events of REM sleep in a patient with severe obstructive sleep apnea syndrome. *J Clin Sleep Med* 2008;4:257–259.

Neurology[®]

Sleep-specific atrioventricular block

Jeffrey M. Ellenbogen

Neurology 2010;74;90

DOI 10.1212/WNL.0b013e3181c7daba

This information is current as of December 28, 2009

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/74/1/90.full>

References

This article cites 3 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/74/1/90.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.