Teaching NeuroImages:

Harlequin syndrome caused by lesion of sympathetic regulatory neurons

Saeed Bohlega, MD,
FAAN
Bent Stigsby, MD, PhD
Fahd Al Mohaileb, MD

Address correspondence and
reprint requests to Dr. Saeed
Bohlega, MBC 76, Department
of Neurosciences, King Faisal
Specialist Hospital and Research
Centre, PO Box 3354, Riyadh
11211, Saudi Arabia
bohlega@kfshrc.edu.sa

A 48-year-old woman presented with a 6-year history of recurrent episodes of exertional asymmetric flushing of the face and head. After exercise, she observed a distinct line of demarcation between the left half of her face, which was red, and the right half, which retained its normal color (figure, A). Decreased right facial temperature and sweating were also noted. These episodes resolved after 1 hour of rest. She had no ptosis or myosis. She had mild trauma to the right neck 2 years prior to this complaint.

MRI of head, neck, and cervicobrachial plexus were normal. CT angiogram of the right internal carotid artery showed a small segmented stenosis (figure, B and C). Loss of flushing on one side of the face

indicates an ipsilateral lesion of sympathetic neurons innervating the face. The absence of Horner syndrome indicates intact oculosympathetic fibers.^{1,2} This rare and clinically striking syndrome may result from occult carotid dissection.³

REFERENCES

1. Drummond PD, Finch PM. Reflex control of facial flushing during body heating in man. *Brain* 1989;112:1351–1358.
2. Wasner G, Maag R, Ludwig J, et al. Harlequin syndrome: one face of many etiologies: case study. *Nat Clin Pract Neurol* 2005;1:54–59.
3. Sarikaya H, Georgiadis D, Baumgartner R. Harlequin syndrome in spontaneous dissection of the cervical carotid artery. *Neurology* 2008;71:1459.

Figure Clinical and radiologic features of harlequin syndrome

(A) The patient after exertion showing loss of right-sided flushing and sweating of the face and head. The left side showed normal flushing. (B) Three-dimensional and (C) 2-dimensional CT angiogram of the carotid artery showing stenosis at the origin of the right internal carotid artery.

From the Departments of Neurosciences and Radiology, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia.

Disclosure: Dr. Bohlega reports no disclosures. Dr. Stigsby serves on the editorial board of *Clinical Neurology and Neurosurgery*. Dr. Al-Mohaileb reports no disclosures.

Neurology®

Teaching *NeuroImages*: Harlequin syndrome caused by lesion of sympathetic regulatory neurons

Saeed Bohlega, Bent Stigsby and Fahd Al Mohaileb

Neurology 2010;74:e106

DOI 10.1212/WNL.0b013e3181e3963c

This information is current as of June 14, 2010

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/74/24/e106.full>

References

This article cites 3 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/74/24/e106.full#ref-list-1>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://n.neurology.org/content/74/24/e106.full##otherarticles>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.