Teaching NeuroImages:

ExtraleSIONal bleeding of conus medullaris cavernoma

N. Montano, MD
F. Signorelli, MD
T. Tufo, MD
S. Sioletic, MD
L. Lauretti, MD
R. Pallini, MD
G. Maira, MD
F. Doglietto, MD

Address correspondence and reprint requests to Dr. Nicola Montano, Institute of Neurosurgery, Catholic University, Largo Agostino Gemelli, 8, 00168 Rome, Italy
nicolamontanomd@yahoo.it

Figure 1 Spinal MRI

Sagittal T2-weighted image showing the conus medullaris cavernoma at time of diagnosis (A); 3 months later (B), extraleSIONal bleeding was evident.

A 42-year-old man with a 9-month history of right leg paresthesias was admitted because of the acute onset of numbness and weakness of the left leg with gait impairment. Neurologic examination demonstrated bilateral lower limb hyporeflexia. MRI showed acute extraleSIONal bleeding of a conus medullaris cavernoma, which had been discovered 3 months before (figure 1, A and B). Surgical removal was performed (figure 2). At 3 months follow-up, the patient had mild lower limb paresthesias; leg weakness had resolved.

Figure 2 Histopathology

Hematoxylin & eosin staining (magnification $\times 100$) showing a large vessel and irregular tissue spaces with cystically enlarged lumina and thin walls filled with hemosiderin pigment, typical features of cavernous hemangioma.

The conus medullaris is a rare location for a cavernous hemangioma.¹ Surgery may be performed in cases with acute, symptomatic hemorrhage, although the benefits remain uncertain.^{1,2}

REFERENCES

1. Zevgaridis D, Medele RJ, Hamburger C, Steiger H-J, Reulen H-J. Cavernous haemangiomas of the spinal cord: a review of 117 cases. *Acta Neurochir* 1999;141:237–245.
2. Kharkar S, Shuck J, Conway J, Rigamonti D. The natural history of conservatively managed symptomatic intramedullary spinal cord cavernomas. *Neurosurgery* 2007;60:865–872.

Neurology®

Teaching *NeuroImages*: Extralesional bleeding of conus medullaris cavernoma

N. Montano, F. Signorelli, T. Tufo, et al.

Neurology 2010;75:e1

DOI 10.1212/WNL.0b013e3181e6209b

This information is current as of July 5, 2010

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/75/1/e1.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/75/1/e1.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord Arteriovenous malformation http://n.neurology.org/cgi/collection/arteriovenous_malformation Clinical neurology history http://n.neurology.org/cgi/collection/clinical_neurology_history MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.