Teaching NeuroImages:

A giant developmental venous anomaly in the absence of a superficial venous drainage system

A.M.G. Fuhler, MD
J.M.C. van Dijk, MD,
PhD
K. Koopman, MD,
PhD
G.J. Luijckx, MD, PhD

Address correspondence and
reprint requests to Dr. A.M.G.
Fuhler, UMC Groningen,
Department of Neurosurgery, PO
Box 30001, 9700 RB Groningen,
the Netherlands
mariskafuhler@hotmail.com

Figure Holohemispheric developmental venous anomaly

(A) T1-weighted MRI with IV gadolinium. The centrum semiovale shows multiple contrast-enhanced linear structures from the cortex to the left lateral ventricle. (B) Digital subtraction angiography. In the absence of a superficial venous drainage system, a holohemispheric developmental venous anomaly is visualized.

A 46-year-old woman presented with new left hemispheric seizures. Enhanced T1 MRI showed a vascular abnormality (figure, A). All other sequences were normal. Cerebral angiography revealed a developmental venous anomaly, with the typical caput medusae appearance (figure, B).

Developmental venous anomalies are generally considered benign variants of the venous system and are thus regarded as symptom-free, although they can be associated with cavernomas which may cause seizures.^{1,2}

In this case, without signs of a cavernoma, the absence of a superficial venous system may be responsible for the seizures, since a fragile equilibrium of inflow and outflow may lead to venous congestion.

REFERENCES

1. Pereira VM, Geibprasert S, Krings T, et al. Pathomechanisms of symptomatic developmental venous anomalies. *Stroke* 2008;39:3201–3215.
2. Ruiz DS, Yilmaz H, Gailloud P. Cerebral venous abnormalities: current concepts. *Ann Neurol* 2009;66:271–283.

From the Departments of Neurosurgery (A.M.G.F., J.M.C.v.D.) and Neurology (K.K., G.J.L.), University Medical Center Groningen, University of Groningen, Groningen, the Netherlands.

Disclosure: The authors report no disclosures.

Neurology®

Teaching *NeuroImages*: A giant developmental venous anomaly in the absence of a superficial venous drainage system

A.M.G. Fuhler, J.M.C. van Dijk, K. Koopman, et al.

Neurology 2010;75:e88

DOI 10.1212/WNL.0b013e3181feb247

This information is current as of November 22, 2010

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/75/21/e88.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/75/21/e88.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Arteriovenous malformation
http://n.neurology.org/cgi/collection/arteriovenous_malformation

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2010 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.