Teaching NeuroImages:

Diffuse posterior leukoencephalopathy in MELAS without stroke-like episodes

Dimitri Renard, MD
Helene Bonnaure, MD
Pierre Labauge, MD,
PhD

Address correspondence and
reprint requests to Dr. Dimitri
Renard, Department of
Neurology, CHU Nîmes, Hôpital
Caremeau, Place du Pr Debré,
30029 Nîmes Cedex 4, France
dimitrirenard@hotmail.com

A 46-year-old woman with a history of seizures, migraine, diabetes mellitus, and neurosensory hearing loss presented with progressive gait unsteadiness. Her mother had the same clinical features. Examination revealed short stature, cognitive deficit, and ataxia. Brain MRI showed diffuse atrophy and posterior leukoencephalopathy (figure). Lactate levels were slightly elevated in both serum and CSF. Mitochondrial DNA analysis revealed the m.3243A>G tRNA^{Leu}(UUR) mutation that causes mitochondrial encephalopathy, lactic acidosis, and stroke-like episodes (MELAS). Brain MRI in MELAS classically shows lesions that predominantly affect the occipital cortex and do not conform to large-vessel territories. The posterior predominance of brain lesions in our

patient, although nonspecific, suggested a MELAS-associated pathology. Diffuse symmetric leukoencephalopathy in patients with the m3243A>G mutation and lacking stroke-like episodes is extremely rare and their pathophysiology is unclear.^{1,2} Diabetes mellitus might also play a role in these radiologic abnormalities.

REFERENCES

1. Apostolova LG, White M, Moore SA, Davis PH. Deep white matter pathologic features in watershed regions: a novel pattern of central nervous system involvement in MELAS. *Arch Neurol* 2005;62:1154–1156.
2. Castillo M, Kwok L, Green C. MELAS syndrome: imaging and proton MR spectroscopic findings. *AJNR Am J Neuroradiol* 1995;16:233–239.

Figure Brain MRI

Brain 3-T MRI showing generalized brain atrophy with severe involvement of the cerebellar vermis on sagittal T1-weighted imaging (A) and diffuse symmetric posterior leukoencephalopathy on axial fluid-attenuated inversion recovery sequences (B–D). Diffusion-weighted imaging and apparent diffusion coefficient sequences were normal. Basal ganglia were normal and there were no intracerebral calcifications on MRI.

From the Departments of Neurology (D.R., P.L.) and Endocrinology (H.B.), CHU Nîmes, Hôpital Caremeau, Nîmes, France.

Disclosure: The authors report no disclosures.

Neurology[®]

Teaching *NeuroImages*: Diffuse posterior leukoencephalopathy in MELAS without stroke-like episodes

Dimitri Renard, Helene Bonnaure and Pierre Labauge

Neurology 2010;75:e9

DOI 10.1212/WNL.0b013e3181e8e73e

This information is current as of July 19, 2010

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/75/3/e9.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/75/3/e9.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
All Cerebrovascular disease/Stroke
http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke
Mitochondrial disorders
http://n.neurology.org/cgi/collection/mitochondrial_disorders
MRI
<http://n.neurology.org/cgi/collection/mri>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.