Teaching *NeuroImages*: Neuroimaging leads to recognition of previously undiagnosed Fabry disease

Laura Fancellu, MD
Giovanni Andrea Deiana,
PhD
GianPietro Sechi, MD

Address correspondence and
reprint requests to Dr. Laura
Fancellu, Department of
Neurosciences, Neurological
Clinic, Viale San Pietro, 10,
07100 Sassari, Italy
laura.fancellu@gmail.com

A 41-year-old man with chronic hypertension presented in a transient, global confusional state. The patient had neuropathic pain and kidney disease, the significance not recognized by his doctors. Family history was positive for hypertension and kidney disease. Brain CT, MRI, and magnetic resonance angiography showed abnormalities typical of Fabry disease (figure).¹ The diagnosis was confirmed by genotype (p.Arg227Gln) and enzymology. Cutaneous and ocular signs indicative for Fabry disease were absent.

For the past year, the patient has been treated with IV α -galactosidase-A, 0.2 mg/Kg every 2 weeks.² His neuropathic pain improved, although

renal function continued to deteriorate. Fabry disease is probably underestimated and diagnosis is frequently delayed.² Typical radiologic manifestations allow diagnosis.¹ Early enzyme replacement therapy reduces the risk of clinical worsening and often ameliorates the prognosis.²

REFERENCES

1. Moore DF, Kaneski CR, Askari H, Schiffmann R. The cerebral vasculopathy of Fabry disease. *J Neurol Sci* 2007; 257:258–263.
2. Mehta A, Beck M, Elliott P, et al. Enzyme replacement therapy with agalsidase alfa in patients with Fabry's disease: an analysis of registry data. *Lancet* 2009;374:1986–1996.

Figure Brain CT, brain MRI, and magnetic resonance angiography

Axial CT scan demonstrated increased attenuation in the pulvinar region (arrows, A), corresponding to the sites of hyperintensity on T1-weighted MRI (the pulvinar sign) (arrows, B). Fluid-attenuated inversion recovery-weighted axial MRI section showed multiple white matter lesions in cerebral hemispheres (C). Magnetic resonance angiography showed basilar dolichoectasia (arrows, D).

From the Department of Neurosciences, Neurological Clinic, University of Sassari, Sassari, Italy.

Disclosure: The authors report no disclosures.

Neurology[®]

Teaching *NeuroImages*: Neuroimaging leads to recognition of previously undiagnosed Fabry disease

Laura Fancellu, Giovanni Andrea Deiana and GianPietro Sechi

Neurology 2010;75;e28

DOI 10.1212/WNL.0b013e3181ec8037

This information is current as of August 9, 2010

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/75/6/e28.full>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/75/6/e28.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
All Cerebrovascular disease/Stroke
http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke
CT
<http://n.neurology.org/cgi/collection/ct>
Metabolic disease (inherited)
http://n.neurology.org/cgi/collection/metabolic_disease_inherited
MRI
<http://n.neurology.org/cgi/collection/mri>
Stroke in young adults
http://n.neurology.org/cgi/collection/stroke_in_young_adults

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.