Dystonic drop foot gait in a patient with manganism

In 1983–1984, a 29-year-old battery factory worker gradually developed gait difficulties and tremor. In 1984, he was found to have severely elevated serum manganese levels and retired from work. Clinical examination in 1984 (videos 1–3 on the *Neurology*® Web site at www.neurology.org) showed bilateral foot dystonia and profound parkinsonism. His gait was shuffling, although he lifted his knees relatively highly. Strength of foot dorsiflexors was normal. He did not respond to dopaminergic drugs and his gait disorder slowly deteriorated to an asymmetric dystonic drop foot gait (videos 4 and 5, recorded in 2008).

Although the walking pattern of this patient with manganism resembled that of a strutting rooster, it was distinct from cock gait as classically described in manganism. Patients with classic cock gait walk on the metatarsophalangeal joints and their heels do not touch the ground.^{1,2}

Jules Janssens, MD, Wim Vandenberghe, MD, PhD, Leuven, Belgium

Address correspondence and reprint requests to Dr. Wim Vandenberghe, Department of Neurology, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium; wim.vandenberghe@uzleuven.be

Disclosure: The authors report no disclosures.

ACKNOWLEDGMENT

W.V. is a Senior Clinical Investigator of the Fund for Scientific Research Flanders (FWO).

1. von Jaksch R. Ueber Mangantoxikosen und Manganophobie. *Münch Med Wochenschr* 1907;20:969–972.
2. Huang CC, Chu NS, Lu CS, Calne DB. Cock gait in manganese intoxication. *Mov Disord* 1997;12:807–808.

Supplemental data at
www.neurology.org

Neurology®

Dystonic drop foot gait in a patient with manganism

Jules Janssens and Wim Vandenberghe

Neurology 2010;75;835

DOI 10.1212/WNL.0b013e3181f0752d

This information is current as of August 30, 2010

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/75/9/835.full>

Supplementary Material

Supplementary material can be found at:
<http://n.neurology.org/content/suppl/2010/08/27/75.9.835.DC1>

References

This article cites 2 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/75/9/835.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Basal ganglia

http://n.neurology.org/cgi/collection/basal_ganglia

Dystonia

<http://n.neurology.org/cgi/collection/dystonia>

Gait disorders/ataxia

http://n.neurology.org/cgi/collection/gait_disorders_ataxia

Other toxicology

http://n.neurology.org/cgi/collection/other_toxicology

Parkinson's disease/Parkinsonism

http://n.neurology.org/cgi/collection/parkinsons_disease_parkinsonism

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.