Teaching NeuroImages:

Unilateral prosopometamorphopsia as a dominant hemisphere-specific disconnection sign

J.-Y. Cho, MD, PhD
S.Y. Moon, MD, PhD
K.-S. Hong, MD, PhD
Y.-J. Cho, MD
S.-C. Kim, MD
S.-I. Hwang, MD
S.-Y. Seo, MD
J.-E. Kim, MD
H.K. Park, MD

Address correspondence and reprint requests to Dr. Hee Kyung Park, Department of Neurology, Ilsan Paik Hospital, Inje University, 2240 Daewha-dong, Ilsanseo-gu, Goyang-city, Gyeonggi-do, 411-706, Korea
jenhkpark@gmail.com

Figure Drawings of facial features as described by the patients, along with the patients' brain MRIs

(A) This figure was drawn based on the first patient's statement. (B) Diffusion-weighted MRI of the first patient demonstrated infarction in the left splenium with a tiny focus on the right side. (C) This figure was drawn according to the second patient's statement. (D) Brain MRI of the second patient showed infarction in the right splenium of the corpus callosum.

We report 2 patients who complained that the left half of faces appeared distorted. A 58-year-old right-handed woman explained that although the left side of the face

was clearly visible, it appeared distorted "like a monster." The left eye looked elongated toward the left ear, while the nose appeared to be bent toward the left cheek, and the mouth toward the chin. She did not have any dyslexia in the left or right visual field. A 53-year-old right-handed man described that the left eyelid of people looked swollen as if they had undergone "failed eyelid surgery," while the nose appeared to be bent downward and the left facial outlines either bulged or writhed. This was the same with pictures of others, but was absent with objects. He did not show hemialexia, color anomia, or optic aphasia. Brain MRIs showed splenic infarctions (figure). The prosopometamorphopsia in the right hemifield was likely caused by the disruption of the pathway from the left occipital area to the right hemisphere. Unilateral prosopometamorphopsia could be a dominant hemisphere-specific disconnection sign in which neurologic abnormalities are observed in the ipsilateral side of the dominant hemisphere.¹ Our cases provide additional evidence that the splenium of the corpus callosum interconnects visual cortices and the right hemisphere is dominant for integrating facial information.²

REFERENCES

1. Lee JI, Kim JH, Lee BH, et al. Dominance specific visual extinction associated with callosal disconnection. *Neurocase* 2010;16:7–14.
2. Minnebusch DA, Daum I. Neuropsychological mechanisms of visual face and body perception. *Neurosci Biobehav Rev* 2009;33:1133–1144.

From the Department of Neurology (J.-Y.C., K.-S.H., Y.-J.C., S.-C.K., S.-I.H., S.-Y.S., J.-E.K., H.K.P.), Ilsan Paik Hospital, Inje University College of Medicine, Gyeonggi-do; and Department of Neurology (S.Y.M.), Ajou University School of Medicine, Suwon, Korea.

Disclosure: Dr. J.-Y. Cho receives research support from the Ministry of Health, Welfare and Family Affairs, Republic of Korea. Dr. Moon reports no disclosures. Dr. Hong serves on the speakers' bureau of Pfizer Inc; and receives/has received research support from Otsuka Pharmaceutical Co., Ltd., Boryung Pharmaceutical Co., Ltd., Novartis, and the Ministry of Health, Welfare and Family Affairs, Republic of Korea. Dr. Y.-J. Cho receives research support from sanofi-aventis and Otsuka Pharmaceutical Co., Ltd. Dr. S.-C. Kim, Dr. Hwang, Dr. Seo, Dr. J.E. Kim, and Dr. Park report no disclosures.

Neurology®

Teaching *NeuroImages*: Unilateral prosopometamorphopsia as a dominant hemisphere-specific disconnection sign

J.-Y. Cho, S.Y. Moon, K.-S. Hong, et al.

Neurology 2011;76:e110

DOI 10.1212/WNL.0b013e31821d74a0

This information is current as of May 30, 2011

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/76/22/e110.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/76/22/e110.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Neuropsychology/Behavior http://n.neurology.org/cgi/collection/all_neuropsychology_behavior Infarction http://n.neurology.org/cgi/collection/infarction MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2011 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.