Teaching NeuroImages:

Isolated bilateral trigeminal nerve palsy

Efthimios Dardiotis, MD
Gianna Patramani, MD
Styliani Ralli, MD
Georgios M.
Hadjigeorgiou, MD

Address correspondence and
reprint requests to Dr. Georgios
M. Hadjigeorgiou, Department of
Neurology, University Hospital of
Larissa, Faculty of Medicine,
University of Thessaly, Biopolis,
Mezourlo Hill, Larissa 41100,
Greece
gmhadji@yahoo.com

Figure 1 Patient with bilateral trigeminal nerve palsy

The patient was unable to close his mouth. Neurologic examination revealed bilateral trigeminal nerve involvement.

A 73-year-old man with a history of non-Hodgkin lymphoma (NHL) presented 6 months after systemic remission was achieved with a painless mandible drop (figure 1), weakness of masseters and pterygoids, loss of corneal reflex, and impaired facial sensation. Brain MRI (figure 2) demonstrated gadolinium enhancement of both trigeminal nerves, and CSF analysis was compatible with NHL recurrence.¹ No other evidence of systemic disease progression was found. The patient experienced partial resolution of his symptoms and signs after administration of intrathecal methotrexate and systemic chemotherapy.

CSF invasion of malignant cells is the most likely mechanism of isolated NHL recurrence in the CNS.²

Figure 2 MRI of the patient

Axial (A) and coronal (B) T2 MRI demonstrating enlargement of both trigeminal nerves (arrows). Axial T1-weighted (C) and postgadolinium images (D) demonstrating gadolinium enhancement of both trigeminal nerves (arrowheads).

However, the biologic mechanism underlying the confined cranial nerve infiltration is still poorly understood, although a neurotropic nature of malignant cells has been hypothesized.³

REFERENCES

1. Chamberlain MC. Leptomeningeal metastasis. *Semin Neurol* 2010;30:236–244.
2. Gleissner B, Chamberlain MC. Neoplastic meningitis. *Lancet Neurol* 2006;5:443–452.
3. Baehring JM, Damek D, Martin EC, Betensky RA, Hochberg FH. Neurolymphomatosis. *Neuro-Oncol* 2003;5:104–115.

Neurology®

Teaching NeuroImages: Isolated bilateral trigeminal nerve palsy

Efthimios Dardiotis, Gianna Patramani, Styliani Ralli, et al.

Neurology 2011;76:e115

DOI 10.1212/WNL.0b013e31821f4622

This information is current as of June 13, 2011

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/76/24/e115.full>

References

This article cites 3 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/76/24/e115.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2011 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.