3. Irani SR, Michell AW, Lang B, et al. Faciobrachial dystonic seizures precede Lgi1 antibody limbic encephalitis. *Ann Neurol* 2011;69:892–900.
4. Irani SR, Buckley C, Vincent A, et al. Immunotherapy-responsive seizure-like episodes with potassium channel antibodies. *Neurology* 2008;71:1647–1648.
5. Vincent A, Buckley C, Schott JM, et al. Potassium channel antibody-associated encephalopathy: a potentially immunotherapy-responsive form of limbic encephalitis. *Brain* 2004;127:701–712.
6. Irani SR, Alexander S, Waters P, et al. Antibodies to Kv1 potassium channel-complex proteins leucine-rich, glioma inactivated 1 protein and contactin-associated protein-2 in limbic encephalitis, Morvan's syndrome and acquired neuromyotonia. *Brain* 2010;133:2734–2748.
7. Mizobuchi M, Matsuda K, Inoue Y, et al. Dystonic posturing associated with putaminal hyperperfusion depicted on subtraction SPECT. *Epilepsia* 2004;45:948–953.
8. Rusu V, Chassoux F, Landré E, et al. Dystonic posturing in seizures of mesial temporal origin: electroclinical and metabolic patterns. *Neurology* 2005;65:1612–1619.
9. Chassagnon S, Namer IJ, Armpach JP, et al. SPM analysis of ictal-interictal SPECT in mesial temporal lobe epilepsy: relationships between ictal semiology and perfusion changes. *Epilepsy Res* 2009;85:252–260.
10. Kuba R, Rektor I, Brázdil M. Ictal limb dystonia in temporal lobe epilepsy: an invasive video-EEG finding. *Eur J Neurol* 2003;10:641–649.
11. Gu W, Brodtkorb E, Steinlein OK. LGI1 is mutated in familial temporal lobe epilepsy characterized by aphasic seizures. *Ann Neurol* 2002;52:364–367.
12. Lai M, Huijbers MG, Lancaster E, et al. Investigation of LGI1 as the antigen in limbic encephalitis previously attributed to potassium channels: a case series. *Lancet Neurol* 2010;9:776–785.

CORRECTION

Sex differences in risk factors for aneurysmal subarachnoid hemorrhage: A cohort study

In the article “Sex differences in risk factors for aneurysmal subarachnoid hemorrhage: A cohort study” by H. Lindekleiv et al. (*Neurology*® 2011;76:637–643), the number of participants from the HUNT cohort was incorrect. The correct number of participants was 65,526 (not 65,580) and the number of aSAH cases during follow-up was 71 (not 69). The corrected results do not change the conclusions published. The authors regret the error.

Neurology[®]

Sex differences in risk factors for aneurysmal subarachnoid hemorrhage: A cohort study

Neurology 2011;77;2143
DOI 10.1212/WNL.0b013e31823ea3aa

This information is current as of December 12, 2011

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/77/24/2143.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2011 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.