Teaching NeuroImages: Sequential MRI of the pituitary in Sheehan syndrome

Ajaykumar Morani, MD
Hemant Parmar, MD
Mohannad Ibrahim,
MD

Correspondence & reprint
requests to Dr. Morani:
ajaycmorani@yahoo.com

Figure 1 Sheehan syndrome

Coronal and sagittal postcontrast T1-weighted spin-echo images show an enlarged pituitary with rim enhancement and no internal enhancement (arrows). Findings are suggestive of nonhemorrhagic pituitary infarct, i.e., Sheehan syndrome.

Figure 2 Temporal evolution of the pituitary gland in Sheehan syndrome

Coronal and sagittal postcontrast T1-weighted spin-echo images show atrophy of the anterior pituitary gland (arrows) on follow-up MRI 1 year later.

A 32-year-old woman with postpartum bleeding was unable to lactate. Investigations revealed hypocortisolism and hypoprolactinemia. MRI revealed an enlarged cystic pituitary suspicious for infarct. Sheehan syndrome is a clinical state of panhypopituitarism resulting from pituitary infarction after a

peripartum hypotensive episode.^{1,2} In the acute phase, the pituitary is enlarged with homogeneous T1-weighted hypointensity and T2-weighted hyperintensity with rim enhancement (figure 1). Sometimes infarcted areas are interspersed with perfused tissue, causing irregular pituitary enhancement.¹ It is later followed by anterior pituitary atrophy (figure 2).^{1,2} The differential is lymphocytic hypophysitis, which may show a similar cystic focus in the pituitary in 5% of patients with subsequent atrophy.¹

AUTHOR CONTRIBUTIONS

Dr. Morani: drafting/revising the manuscript, study concept or design, acquisition of data. Dr. Parmar: drafting/revising the manuscript, analysis or interpretation of data, study supervision. Dr. Ibrahim: study concept or design, acquisition of data.

REFERENCES

1. Kaplun J, Fratila C, Ferenczi A, et al. Sequential pituitary MR imaging in Sheehan syndrome: report of 2 cases. *AJNR* 2008;29(suppl 5):941–943.
2. Zak IT, Dulai HS, Kish KK. Imaging of neurologic disorders associated with pregnancy and the postpartum period. *Radiographics* 2007;27(suppl 1):95–108.

From the Department of Radiology, University of Michigan, Ann Arbor.

Disclosure: Dr. Morani reports no disclosures. Dr. Parmar serves as an Associate Editor for *Pediatric Neuroradiology*. Dr. Ibrahim reports no disclosures.

Neurology®

Teaching *NeuroImages*: Sequential MRI of the pituitary in Sheehan syndrome

Ajaykumar Morani, Hemant Parmar and Mohannad Ibrahim

Neurology 2012;78:e3

DOI 10.1212/WNL.0b013e31823ed07d

This information is current as of December 26, 2011

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/78/1/e3.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/78/1/e3.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2012 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.