Basilar leptomeningitis in Vogt-Koyanagi-Harada disease

Figure 1 Optic coherence tomography (OCT) and funduscopy

OCT and funduscopy on presentation (left) and following steroids (right). OCT (A) demonstrates serous exudate detaching the retina (arrow) from the pigmented layer (arrowhead) with normalization after steroids (B). The left (C) and right (E) fundi show retinal detachments (black arrowheads) and papillitis, which resolve with steroids, (D) and (F).

A 35-year-old Filipino man had acute visual loss, tinnitus, dysacusis, and severe horizontal vertigo. Examination revealed bidirectional gaze-evoked nystagmus and panuveitis with bilateral inferior serous retinal detachments (figure 1) characteristic of Vogt-Koyanagi-Harada disease.¹ Brain MRI demonstrated mesencephalic, pontine, and cerebellar leptomeningeal enhancement (figure 2). There was cerebrospinal lymphocytic pleocytosis. Following IV Solu-Medrol, his fundus was significantly improved.

Vogt-Koyanagi-Harada disease is a uveomeningeal syndrome of panuveitis with bullous serous retinal detachment, meningitis, and dysacusis.¹ Although the targets of inflammation, melanin-containing cells,^{1,2} are predominantly located over the ventral medulla,³ our patient demonstrated a ponto-mesencephalic and cerebellar predilection of inflammation.

Figure 2 Gadolinium-enhanced MRI of the brain

(A) Axial, (B) coronal, and (C) sagittal gadolinium-enhanced magnetic resonance brain images demonstrate leptomeningeal enhancement of the midbrain (arrow), ventral pons (outline arrow), and cerebellum (arrowheads).

Yince Loh, MD, Tacoma, WA

Disclosure: The author reports no disclosures.

The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Army, the Department of Defense, or the United States Government.

Correspondence & reprint requests to Dr. Loh: yincer@yahoo.com

1. Brazis PW, Stewart M, Lee AG. The uveo-meningeal syndromes. *Neurologist* 2004;10:171–184.
2. Reed RW, Rao NA, Cunningham ET Jr. Vogt-Koyanagi-Harada disease. *Curr Opin Ophthalmol* 2000;11:437–442.
3. Goldgeier MH, Klein LE, Klein-Angerer S, Moellmann G, Nordlund JJ. The distribution of melanocytes in the leptomeninges of the human brain. *J Invest Dermatol* 1984;82:235–238.

Neurology®

Basilar leptomeningitis in Vogt-Koyanagi-Harada disease

Yince Loh

Neurology 2012;78;438-439

DOI 10.1212/WNL.0b013e318245d2e3

This information is current as of February 6, 2012

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/78/6/438.full>

References

This article cites 3 articles, 0 of which you can access for free at:
<http://n.neurology.org/content/78/6/438.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Autoimmune diseases

http://n.neurology.org/cgi/collection/autoimmune_diseases

Meningitis

<http://n.neurology.org/cgi/collection/meningitis>

MRI

<http://n.neurology.org/cgi/collection/mri>

Retina

<http://n.neurology.org/cgi/collection/retina>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:

<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2012 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.