Teaching *NeuroImages*: Crossed cerebellar diaschisis in hemispheric status epilepticus

Allie M. Massaro, MD

Correspondence & reprint
requests to Dr. Massaro:
allie.massaro@uphs.upenn.edu

Figure 1 Continuous EEG at 15 $\mu\text{V/mm}$ illustrating right frontotemporal seizure activity characterized by rhythmic theta/delta admixed with sharp waves over the right frontotemporal region

Figure 2 Axial brain MRI diffusion-weighted imaging (A) and corresponding axial brain MRI apparent diffusion coefficient (B)

Axial brain MRI diffusion-weighted imaging sequence demonstrates restrictive diffusion of the right cortex, right basal ganglia, and left cerebellum (A). Corresponding apparent diffusion coefficient mapping shows restricted diffusion in the right cortex (B).

A 36-year-old woman with Sheehan syndrome was found unresponsive. Evaluation was consistent with Addisonian crisis and myxedema coma. Continuous EEG demonstrated right frontotemporal nonconvulsive status initially refractory to multiple antiepileptic drugs (figure 1).

MRI brain revealed restricted diffusion (figure 2) within the cortex of the right hemisphere and left cerebellum, suggestive of focal status epilepticus. There was no associated enhancement on apparent diffusion coefficient (ADC) (figure 2), and near normalization of ADC mapping on follow-up imaging. Although primarily

recognized in the stroke literature, crossed cerebellar diaschisis may represent injury caused by excessive neuronal transmission from prolonged excitatory synaptic activity via the cortico-pontine-cerebellar pathways.^{1,2}

REFERENCES

1. Samaniego EA, Stuckert E, Fischbein N, Wijman CA. Crossed cerebellar diaschisis in status epilepticus. *Neurocrit Care* 2010; 12:88–90.
2. Tien RD, Ashdown BC. Crossed cerebellar diaschisis and crossed cerebellar atrophy: correlation of MR findings, clinical symptoms, and supratentorial diseases in 26 patients. *AJR Am J Roentgenol* 1992;158:1155–1159.

From the Department of Neurology, Hospital of the University of Pennsylvania, University of Pennsylvania, Philadelphia.
The author reports no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Neurology®

Teaching *NeuroImages*: Crossed cerebellar diaschisis in hemispheric status epilepticus

Allie M. Massaro

Neurology 2012;79:e182

DOI 10.1212/WNL.0b013e318274de51

This information is current as of November 12, 2012

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/79/20/e182.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/79/20/e182.full#ref-list-1
Citations	This article has been cited by 2 HighWire-hosted articles: http://n.neurology.org/content/79/20/e182.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Coma http://n.neurology.org/cgi/collection/coma DWI http://n.neurology.org/cgi/collection/dwi EEG http://n.neurology.org/cgi/collection/eeg_ MRI http://n.neurology.org/cgi/collection/mri Status epilepticus http://n.neurology.org/cgi/collection/status_epilepticus
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2012 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.