Teaching *NeuroImages*: Isolated hypoglossal nerve palsy due to internal carotid artery dissection

Taro Okunomiya, MD
Takashi Kageyama, MD,
PhD
Toshihiko Suenaga, MD,
PhD

Correspondence & reprint
requests to Dr. Suenaga:
suenaga@tenriyorozeu.jp

Figure Photograph and brain MRI

(A) The patient's tongue deviated to the left. (B) T1-weighted imaging demonstrates narrowing of the lumen and intramural hematoma in the left internal carotid artery. (C) True fast imaging with steady-state precession reveals the anatomic juxtaposition of the hypoglossal nerve (arrowhead) and the dissected internal carotid artery (arrow).

A 57-year-old man presented with dysarthria and left-sided headache lasting 4 days. Neurologic examination revealed deviation of the tongue to the left (figure, A). MRI demonstrated the left carotid artery dissection (figure, B) and its anatomic juxtaposition with the left hypoglossal nerve (figure, C). Tongue deviation returned to normal after 4 weeks of treatment with aspirin.

Acute tongue deviation with headache is an important manifestation of internal carotid artery dissection. The hypothesized mechanism is direct compression of the hypoglossal nerve by the expanded dissected arterial

wall. Carotid artery dissection affected the hypoglossal nerve in 5% of cases.¹

AUTHOR CONTRIBUTIONS

T. Okunomiya: study concept and design, drafting and revising the manuscript, analysis or interpretation of data. T. Kageyama: critically revising the manuscript for intellectual content. T. Suenaga: study concept and design, drafting and revising the manuscript, analysis or interpretation of data, guarantor of study.

REFERENCE

1. Mokri B, Silbert PL, Schievink WI, Piepgras DG. Cranial nerve palsy in spontaneous dissection of the extracranial internal carotid artery. *Neurology* 1996;46:356–359.

From the Department of Neurology, Tenri Hospital, Tenri City, Japan.

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Neurology®

Teaching *NeuroImages*: Isolated hypoglossal nerve palsy due to internal carotid artery dissection

Taro Okunomiya, Takashi Kageyama and Toshihiko Suenaga

Neurology 2012;79:e37

DOI 10.1212/WNL.0b013e3182604506

This information is current as of July 23, 2012

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/79/4/e37.full
References	This article cites 1 articles, 1 of which you can access for free at: http://n.neurology.org/content/79/4/e37.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Carotid artery dissection http://n.neurology.org/cgi/collection/carotid_artery_dissection MRI http://n.neurology.org/cgi/collection/mri Secondary headache disorders http://n.neurology.org/cgi/collection/secondary_headache_disorders
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2012 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.