brain infarcts and white matter lesions increase stroke risk in the general population: the Rotterdam Scan Study. *Stroke* 2003;34:1126–1129.

3. Cupini LM, Diomedes M, Placidi F, Silvestrini M, Giacomini P. Cerebrovascular reactivity and Subcortical infarctions. *Arch Neurol* 2001;58:577–581.
4. Silvestrini M, Viticchi G, Falsetti L, et al. The role of carotid atherosclerosis in Alzheimer's disease progression. *J Alzheimers Dis* 2011;25:719–726.
5. Wahlund LO, Barkhof F, Fazekas F, et al. A new rating scale for age-related white matter changes applicable to MRI and CT. *Stroke* 2001;32:1318–1322.

A COMPARISON OF TAU AND 14-3-3 PROTEIN IN THE DIAGNOSIS OF CREUTZFELDT-JAKOB DISEASE

Andre Karch, Inga Zerr, Göttingen, Germany:

Hamlin et al.¹ compared validities of 2 commonly used CSF tests for sporadic Creutzfeldt-Jakob disease (sCJD)—14-3-3 and total tau—and presented results contrary to previous studies.^{1–4} The present study is restricted to autopsy-confirmed cases, which seems reasonable, since neuropathology is the diagnostic gold standard in sCJD. However, using only autopsy-confirmed cases can also cause severe selection bias in a prospective setting, given that these tests are frequently requested in potentially treatable conditions.^{2,3} In the present study, clinicians were given results of 14-3-3 but not tau tests during the patient's lifetime. Autopsies were obtained in fewer than 10% of all initially referred patients. Thereby, decision about initiation of

postmortem autopsy was directly dependent on 14-3-3 but not tau results. It seems a fair assumption that patients with CJD and negative 14-3-3 as well as patients without CJD and positive 14-3-3 were more likely to be autopsied than patients with consistent clinical history and 14-3-3 test results. Thereby, specificity and sensitivity of 14-3-3 are underestimated in this study design, which could explain differences in 14-3-3 validity when compared to previous studies. A final decision on test accuracy of 14-3-3 and total tau needs careful assessment in a prospective, multicenter study using standardized protocols for laboratory, clinical, and pathologic data.⁵

© 2013 American Academy of Neurology

1. Hamlin C, Puoti G, Berri S, et al. A comparison of tau and 14-3-3 protein in the diagnosis of Creutzfeldt-Jakob disease. *Neurology* 2012;79:547–552.
2. Chitravas N, Jung R, et al. Treatable neurological disorders misdiagnosed as Creutzfeldt-Jakob disease. *Ann Neurol* 2011;70:437–444.
3. Stoeck K, Sanchez-Juan P, Gawinecka J, et al. CSF biomarker supported diagnosis of CJD and rapid dementias: a longitudinal multicenter study over 10 years. *Brain* 2012; 135:3051–3061.
4. Sanchez-Juan P, Green A, Ladogana A, et al. CSF tests in the differential diagnosis of Creutzfeldt-Jakob disease. *Neurology* 2006;67:637–643.
5. DEMTEST (biomarker-based diagnosis of rapid progressive dementias: optimisation of diagnostic protocols). Available at: <http://www.neurodegenerationresearch.eu/initiatives/biomarker-transnational-call/results-of-funding-call/demtest/>. Accessed April 9, 2013.

CORRECTIONS

Teaching *NeuroImages*: Basal ganglia involvement in facio-brachial dystonic seizures associated with LGI1 antibodies

In the article “Teaching *NeuroImages*: Basal ganglia involvement in facio-brachial dystonic seizures associated with LGI1 antibodies” by D. Plantone et al. (*Neurology*® 2013;80:e183–e184), figures 1 and 2 were transposed and should be reversed. The publisher regrets the error.

Severe congenital *RYR1*-associated myopathy: The expanding clinicopathologic and genetic spectrum

In the article “Severe congenital *RYR1*-associated myopathy: The expanding clinicopathologic and genetic spectrum” by D. Bharucha-Goebel et al. (*Neurology*® 2013;80:1584–1589), there is an error in the author list. The fourth author should have read Kristen Zukosky. The editorial staff regrets the error.

Author disclosures are available upon request (journal@neurology.org).

Neurology®

Teaching *NeuroImages*: Basal ganglia involvement in facio-brachial dystonic seizures associated with LGI1 antibodies

Neurology 2013;80;2081
DOI 10.1212/WNL.0b013e31829926a2

This information is current as of May 27, 2013

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/80/22/2081.2.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.