Editors' Note: In response to "Hemichorea-hemiballism associated with hyperglycemia and a developmental venous anomaly" by Kalia et al., Drs. Civardi and Collini describe their own case of putaminal CT hyperperfusion without the developmental venous anomaly (DVA), resulting in unilateral myoclonus. The authors describe possible explanations for the abnormal CT perfusion (CTP) findings in their own case study.

Megan Alcauskas, MD, and Robert C. Griggs, MD

HEMICHOREA-HEMIBALLISM ASSOCIATED WITH HYPERGLYCEMIA AND A DEVELOPMENTAL VENOUS ANOMALY

Carlo Civardi, Alessandra Collini, Novara, Italy:

Kalia et al.¹ described a case of a hyperglycemia-induced hemichorea-hemiballism with the typical high signal on T1-weighted images on MRI² in the putamen contralateral to the movement. Additional neuroimaging revealed a DVA adjacent to the affected putamen associated with increased cerebral blood flow and volume on CTP.

The authors proposed that altered hemodynamics within the basal ganglia along with a metabolic disturbance resulted in movement disorder. We reported a man with repetitive myoclonus of the left side of the body and contralateral putaminal CT hyperperfusion. Carbamazepine completely controlled myoclonus and, concurrently, CT showed normal perfusion.³ Our patient did not have a DVA and there was no metabolic defect.

As in other hyperkinetic disorders, CT hyperperfusion as observed in these patients^{1,3} may indicate failed autoregulatory mechanisms in the basal ganglia vessels.⁴ These 2 reports showed the same putaminal hyperperfusion, albeit with a different etiology. The perfusional pattern may be related to hyperkinetic status and not to the etiology of the movement disorder. Moreover, this recent report confirms the utility of CTP in the assessment of hyperkinetic disorders.

Author Response: Lorraine V. Kalia, Anthony E. Lang, Richard I. Aviv, Mario Masellis, Toronto: We thank Drs. Civardi and Collini for their thoughtful comments. We agree that there are several potential explanations for the abnormal CTP findings in our

case.¹ As they suggested, one possibility is that the observed increases in cerebral blood flow and cerebral blood volume are a consequence and not a cause of the patient's hemiballism-hemichorea. Another possibility is that this CTP pattern within the putamen was a result of the adjacent DVA. A similar pattern has been previously described with MRI perfusion (MRP) in 4 patients with a DVA.⁵ Follow-up imaging in our patient after complete resolution of her hemiballism-hemichorea may have been useful in differentiating between these 2 possibilities (i.e., resolution of CTP abnormalities with the first possibility vs persistence with the second possibility). However, based on similarities between the MRP and CTP techniques, we anticipate that the latter would be more likely. While the CTP findings in our case¹ and those described by Civardi et al.³ are interesting, further characterization of this type of imaging in hyperkinetic movement disorders is required before we can establish its utility in the assessment of patients with movement disorders.

© 2013 American Academy of Neurology

1. Kalia LV, Mozessohn L, Aviv RI, et al. Hemichorea-hemiballism associated with hyperglycemia and a developmental venous anomaly. *Neurology* 2012;78:838–839.
2. Postuma RB, Lang AE. Hemiballism: revisiting a classic disorder. *Lancet Neurol* 2003;2:661–668.
3. Civardi C, Collini A, Stecco A, Carriero A, Monaco F. Neurological picture: putamen hyperperfusion in subcortical-supraspinal myoclonus. *J Neurol Neurosurg Psychiatry* 2010;81:330.
4. Hsu JL, Wang HC, Hsu WC. Hyperglycemia-induced unilateral basal ganglion lesions with and without hemichorea: a PET study. *J Neurol* 2004;251:1486–1490.
5. Camacho DL, Smith JK, Grimme JD, Keyserling HF, Castillo M. Atypical MR imaging perfusion in developmental venous anomalies. *AJNR Am J Neuroradiol* 2004;25:1549–1552.

DECREASED IRON LEVELS IN THE TEMPORAL CORTEX IN POSTMORTEM HUMAN BRAINS WITH PARKINSON DISEASE

Osamu Kano, Ken Ikeda, Yasuo Iwasaki, Tokyo:

Yu et al.¹ reported that iron levels in the temporal cortex were reduced in patients with Parkinson disease (PD) compared with age-matched controls. They also determined that patients with Alzheimer disease (AD) had no change in iron levels in the temporal cortex. We

would like to know whether Yu et al.¹ included any patients with memory disturbance in this study.

Sofic et al.² reported an increase in iron levels in the substantia nigra of patients with PD, and showed a decrease in iron levels in the temporal cortex, thereby suggesting that the increase in iron levels may be associated with parkinsonian symptoms. In addition, did Yu et al. find a relationship between iron level in the temporal cortex and cognitive disability scales used in their study? Symptoms associated with the temporal lobe are not seen in patients with typical PD, so we would like to know the iron levels in other brain regions.

Iron may be associated with pathogenesis in PD and further study is needed to determine the relationship between iron and its distribution.

Author Response: Hong Jiang, Qingdao, China: Memory disturbances were present in patients with AD in this study. However, no changes were observed in patients with PD, so we did not find a relationship between iron levels in the temporal cortex and cognitive disability scales in our cohort. Iron levels in other brain regions were not determined because our access to samples was restricted.

© 2013 American Academy of Neurology

1. Yu X, Du T, Song N, et al. Decreased iron levels in the temporal cortex in postmortem human brains with Parkinson disease. *Neurology* 2013;80:492–495.
2. Sofic E, Riederer P, Heinsen H, et al. Increased iron (III) and total iron content in post mortem substantia nigra of parkinsonian brain. *J Neural Transm* 1988;74: 199–205.

***Neurology*[®] WriteClick: Join the Debate!**

The editors encourage comments about recent articles through WriteClick:

Go to www.neurology.org and click on the “WriteClick” tab at the top of the page. Responses will be posted within 72 hours of submission.

Before using WriteClick, remember the following:

- WriteClick is restricted to comments about studies published in *Neurology* within the last eight weeks
- Read previously posted comments; redundant comments will not be posted
- Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
- You can include a maximum of five authors (including yourself)

Author disclosures are available upon request (journal@neurology.org).

Neurology®

Decreased iron levels in the temporal cortex in postmortem human brains with Parkinson disease

Osamu Kano, Hong Jiang, Ken Ikeda, et al.

Neurology 2013;81;1181-1182

DOI 10.1212/01.wnl.0000435455.95141.16

This information is current as of September 23, 2013

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/81/13/1181.2.full>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/81/13/1181.2.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.