Teaching *NeuroImages*: Snake eyes appearance in MRI in patient with ALS

Sushma Sharma, MD
Aditya Murgai, MD
Pradeep Pankajakshan
Nair, MD, DM
Ananthakrishnan
Ramesh, MD

Correspondence to
Dr. Nair:
drpradeepnair17@gmail.com

Figure MRI cervical spine

(A) MRI cervical spine axial T1-weighted sequence shows symmetrical hypointense signals in the anterior horn. (B) MRI cervical spine axial T2-weighted sequence shows symmetrical hyperintense signals in the anterior horn.

A 17-year-old girl presented with 6 months of progressive asymmetrical distal weakness of upper limbs with fasciculation. There were no bulbar symptoms, weakness in lower limbs, or sensory symptoms. There was no history of neck injury or pain. Deep tendon reflexes were all brisk. MRI of cervical spine revealed symmetrical T1 hypointense, T2 hyperintense signal in the anterior horns ("snake eyes" appearance) (figure, A and B) without evidence of extradural compression. EMG showed diffuse denervation changes in both upper and lower limb muscles. Sensory nerve action potentials were normal. A diagnosis of amyotrophic lateral sclerosis (ALS) was made. Snake eyes appearance has been described in disorders like ALS, cervical spondylotic amyotrophy, Hirayama disease, and ossification of posterior longitudinal ligament.^{1,2}

AUTHOR CONTRIBUTIONS

Sushma Sharma: data acquisition, drafting of manuscript. Aditya Murgai: data acquisition, drafting of manuscript. Pradeep Pankajakshan

Nair: drafting and revising of manuscript, concept of the manuscript. Ananthakrishnan Ramesh: data acquisition and revising of manuscript.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

REFERENCES

1. Friedman DP, Tartaglino LM, Fisher AR, Flanders AE. MR imaging in the diagnosis of intramedullary spinal cord diseases that involve specific neural pathways or vascular territories. *AJR Am J Roentgenol* 1995;165: 515–523.
2. Mizuno J, Nakagawa H, Inoue T, Hashizume Y. Clinico-pathological study of "snake-eye appearance" in compressive myelopathy of the cervical spinal cord. *J Neurosurg* 2003; 99(2 suppl):162–168.

Neurology®

Teaching *NeuroImages*: Snake eyes appearance in MRI in patient with ALS

Sushma Sharma, Aditya Murgai, Pradeep Pankajakshan Nair, et al.

Neurology 2013;81:e29

DOI 10.1212/WNL.0b013e31829d85ae

This information is current as of July 29, 2013

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/81/5/e29.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2013/07/28/81.5.e29.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/81/5/e29.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/81/5/e29.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord Amyotrophic lateral sclerosis http://n.neurology.org/cgi/collection/amyotrophic_lateral_sclerosis_MRI MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.