Reversible leukoencephalopathy in sodium monofluoroacetate intoxication

Figure Serial diffusion-weighted imaging and magnetic resonance spectroscopy findings

Initial diffusion-weighted imaging (DWI) shows restricted diffusion in widespread white matter (A) and magnetic resonance spectroscopy (^1H -MRS) shows normal *N*-acetylaspartate (NAA)/creatine and choline/creatine ratio (B). Follow-up DWI and ^1H -MRS performed 1 month later reveal complete resolution of white matter abnormalities (C) with unchanged NAA/creatine and choline/creatine ratio (D).

A 38-year-old man presented with vomiting and progressive mental deterioration after ingesting 210 mL sodium monofluoroacetate solution (SMFA, 2%), a toxic rodenticide, in a suicide attempt. Neither seizure nor tetany was observed. Laboratory tests including ECG, arterial blood gas, serum calcium, and electrolyte were unremarkable. Initial MRI revealed extensive white matter restricted diffusion without disrupted neuronal integrity (figure, A and B). He improved to normal over 2 weeks. Follow-up MRI showed resolution of white matter abnormalities with unaltered metabolic markers (figure, C and D). SMFA selectively inhibits glial Krebs cycle and lowers energy production, with eventual cell death.¹ Reversible MRI cytotoxic edema without neuronal loss² suggests that SMFA intoxication is in the differential diagnosis of reversible leukoencephalopathy.

Jung Bin Kim, MD, Ji-Wan Jang, MD, Ji Hyun Kim, MD, PhD

From Korea University Guro Hospital, Korea University College of Medicine, Seoul.

Author contributions: J.B. Kim and Dr. Jang contributed to preparation of the images and drafting the manuscript. J.H. Kim contributed to supervision of the study and revising the manuscript.

Study funding: No targeted funding reported.

Disclosure: The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

Correspondence to Dr. J.H. Kim: jhkim.merrf@gmail.com

1. Hassel B, Paulsen RE, Johnsen A, Fonnum F. Selective inhibition of glial cell metabolism in vivo by fluorocitrate. *Brain Res* 1992;576:120–124.
2. Im TH, Yi HJ. The utility of early brain imaging (diffusion magnetic resonance and single photon emission computed tomography scan) in assessing the course of acute sodium monofluoroacetate intoxication. *J Trauma* 2009;66:E72–E74.

Neurology®

Reversible leukoencephalopathy in sodium monofluoroacetate intoxication

Jung Bin Kim, Ji-Wan Jang and Ji Hyun Kim

Neurology 2014;82;1190-1191

DOI 10.1212/WNL.0000000000000262

This information is current as of March 31, 2014

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/82/13/1190.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/82/13/1190.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): DWI http://n.neurology.org/cgi/collection/dwi MRI http://n.neurology.org/cgi/collection/mri MRS http://n.neurology.org/cgi/collection/mrs Other toxicology http://n.neurology.org/cgi/collection/other_toxicology
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.