deficit/hyperactivity disorder (ADHD) symptoms in adults is a risk factor for DLB and not for Alzheimer disease.² It is striking that these symptoms were not mentioned in the article by Boot et al.² since our results need confirmation. Regardless, we think that the greater likelihood of having a history of depression or anxiety in patients with diagnosis of DLB would support our findings because the diagnosis of ADHD is often confused or overlaps with depression and anxiety. Comorbidity between these entities and the possibility of bias is well known.^{3–5} In this study, the possibility of such bias is heightened because the data for depression or anxiety were generated solely from medical history and ADHD is usually underreported.

© 2014 American Academy of Neurology

1. Boot BP, Orr CF, Ahlskog JE, et al. Risk factors for dementia with Lewy bodies: a case-control study. *Neurology* 2013; 81:1–8.
2. Golimstok A, Rojas JI, Romano M, Zurru MC, Doctorovich D, Cristiano E. Previous adult attention-deficit and hyperactivity disorder symptoms and risk of dementia with Lewy bodies: a case-control study. *Eur J Neurol* 2011;18:78–84.
3. Michielsen M, Comijs HC, Semeijn EJ, Beekman AT, Deeg DJ, Sandra Kooij JJ. The comorbidity of anxiety and depressive symptoms in older adults with attention-deficit/hyperactivity disorder: a longitudinal study. *J Affect Disord* 2013;148:220–227.
4. Barkley RA, Brown TE. Unrecognized attention-deficit/hyperactivity disorder in adults presenting with other psychiatric disorders. *CNS Spectr* 2008;13:977–984.
5. Chao CY, Gau SS, Mao WC, Shyu JF, Chen YC, Yeh CB. Relationship of attention-deficit-hyperactivity disorder symptoms, depressive/anxiety symptoms, and life quality in young men. *Psychiatry Clin Neurosci* 2008;62:421–426.

INCIDENCE AND PREVALENCE OF SMALL-FIBER NEUROPATHY: A SURVEY IN THE NETHERLANDS

Peter James Dyck, Rochester, MN: Peters et al.¹ described the prevalence and incidence of “pure

small-fiber sensory neuropathy” in a Dutch population. The data appear to be important but the criteria for the disease conditions being tallied remain unclear.

The authors stated: “SFN was diagnosed based on the presence of at least 2 of the following symptoms not otherwise explained: neuropathic pain (burning, shooting, or itching), sheet or sock intolerance, restless legs syndrome, autonomic dysfunction (Sicca syndrome, accommodation problems, hyperhidrosis or hypohidrosis, micturition disturbances, impotence or diminished ejaculation or lubrication, bowel disturbances, hot flushes, orthostatic dizziness, cardiac palpitations), and clinical signs of small-fiber damage (e.g., pinprick loss, thermal sensory loss, allodynia, or hyperalgesia), normal nerve conduction study, and reduced IENFD at the ankle or abnormal quantitative sensory testing thermal thresholds at the foot.”

Since small-fiber sensory and autonomic neuropathy may be asymptomatic, were these cases excluded from this diagnosis? In addition, were known causes (genetic mutations, autoimmune, metabolic, and other known causes) included or excluded? Perhaps the greatest concern is lack of specificity of symptoms and neuropathic signs and tests. Symptoms like palpitation, itching, and Sicca occur more commonly from other diseases than from small-fiber polyneuropathy.

Many of these symptoms are related to age, so were old age–related symptoms included? What specific criteria were used for signs and nerve tests? Descriptors such as abnormal nerve conduction are not specific enough: which attributes and what percentile abnormality? Providing more specific inclusion and exclusion criteria will make the data more useful.

© 2014 American Academy of Neurology

1. Peters MJH, Bakkers M, Merckies ISJ, Hoeijmakers JGJ, van Raak EPM, Faber CG. Incidence and prevalence of small-fiber neuropathy: a survey in the Netherlands. *Neurology* 2013;81:1356–1360.

WriteClick: Rapid Online Correspondence

Have a comment on a recent *Neurology*[®] article you would like to share? Now it is easier and more convenient. *Neurology.org* has launched WriteClick on the home page and sidebars of each article to encourage remarks and debate among users.

WriteClick is restricted to comments about studies published in *Neurology* within the last eight weeks.

Learn more at <http://www.neurology.org/letters>

Author disclosures are available upon request (journal@neurology.org).

Neurology®

Incidence and prevalence of small-fiber neuropathy: A survey in the Netherlands

Peter James Dyck

Neurology 2014;82;1385

DOI 10.1212/01.wnl.0000446502.87636.91

This information is current as of April 14, 2014

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/82/15/1385.full>

References

This article cites 1 articles, 1 of which you can access for free at:
<http://n.neurology.org/content/82/15/1385.full#ref-list-1>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.