Mystery Case: Eyelid myoclonia with absences in an adult patient

Figure 1 Eyelid myoclonia without absence

With levetiracetam: posterior-predominant spiky alpha frequency activity that rapidly spreads to the frontal regions coinciding with eyelid myoclonia.

A 28-year-old man presented to the epilepsy monitoring unit (EMU) with frequent “eye fluttering” episodes since he was 3 years old (video on the *Neurology*[®] Web site at www.neurology.org). He was diagnosed with epilepsy as a teenager after he developed generalized convulsions at age 12. His convulsions were well-controlled with antiepileptic drug therapy. His neurologic examination was normal.

His EMU study revealed brief episodes of eyelid myoclonia (video) coinciding with a paroxysmal “spiky” posterior alpha activity, which rapidly spread to the frontal regions while he was on levetiracetam (figure 1). These episodes were more prominent in light compared to dark and were often triggered by eye closure. Interictally, the posterior dominant rhythm appeared sharply contoured. Discontinuation of levetiracetam for 3 days resulted in 2 to 3 Hz generalized polyspike and wave activity associated with some of the episodes of eyelid myoclonia, within seconds of eyelid closure (figure 2). Additionally, in sleep there

were fragmentary bursts of anteriorly predominant 3 Hz abortive spike and wave discharges. Prior records indicated a photoparoxysmal EEG response.

The patient’s clinical and electrographic picture is consistent with eyelid myoclonia with absences (EMA), also known as Jeavons syndrome.

EMA is an underrecognized syndrome of unknown etiology defined by the triad of childhood onset, photosensitivity, and eyelid myoclonia with or without absence seizures.^{1,2} Patients often develop generalized tonic-clonic seizures in adolescence.^{1,3} The photosensitivity component may decrease as the patients get older.^{1,4} Classic EEG features include a sharply contoured or spiky alpha frequency activity that rapidly spreads to the frontal regions with sustained eye closure. Eyelid myoclonia often coincides as well with occipital epileptic discharges or occipital polyspike and wave discharges, with or without generalized polyspike and wave discharges.³ The overall prognosis of this syndrome is often good as the generalized tonic-clonic and absence

EEG recording showing generalized spike and wave activity. The recording includes channels for various electrodes: Fp1-F7, F7-T7, T7-P7, P7-O1, Fp2-F8, F8-T8, T8-P8, P8-O2, Fp1-F9, F9-C3, C3-P3, P3-O1, Fp2-F8, F8-C4, C4-P4, P4-O2, Fp-Cz, Cz-Pc, and LOG-ROC. The recording is divided into two sections: 'Eyelid myoclonia' and 'Generalized spike and wave'. A scale bar indicates 1000 ms and 70 μ V.

seizures become controlled with antiepileptic drug therapy. Despite the sensitivity of the absences to antiepileptic drugs, the eyelid myoclonia often persists, as seen in our patient.⁴ A small proportion of patients may continue to have uncontrolled generalized convulsions.⁴ Valproic acid, lamotrigine, and levetiracetam are good treatment options,^{1,5} whereas sodium channel agents such as carbamazepine may exacerbate the seizures.⁶ Our patient initially responded to valproic acid, but he developed a reaction concerning for Stevens-Johnson syndrome. With levetiracetam treatment, his generalized tonic-clonic and absence seizures resolved.

Most adult patients with EMA are diagnosed in childhood, but making the diagnosis in adults for the first time may be challenging. Neurology residents and fellows should be aware of the characteristic clinical and electrographic features of this underrecognized syndrome.

Dr. Hannawi analyzed and interpreted the data, drafted and revised the manuscript. Dr. Satpute analyzed and interpreted the data. Dr. Maheshwari analyzed and interpreted the data, reviewed and revised the manuscript.

No targeted funding reported.

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

1. Striano S, Capovilla G, Sofia V, et al. Eyelid myoclonia with absences (Jeavons syndrome): a well-defined idiopathic generalized epilepsy syndrome or a spectrum of photosensitive conditions? *Epilepsia* 2009;50(suppl 5):15–19.
2. Striano S, Striano P, Nocerino C, et al. Eyelid myoclonia with absences: an overlooked epileptic syndrome? *Neurophysiol Clin* 2002;32:287–296.
3. Viravan S, Go C, Ochi A, Akiyama T, Carter Snead O III, Otsubo H. Jeavons syndrome existing as occipital cortex initiating generalized epilepsy. *Epilepsia* 2011; 52:1273–1279.
4. Giannakodimos S, Panayiotopoulos CP. Eyelid myoclonia with absences in adults: a clinical and video-EEG study. *Epilepsia* 1996;37:36–44.
5. Striano P, Sofia V, Capovilla G, et al. A pilot trial of levetiracetam in eyelid myoclonia with absences (Jeavons syndrome). *Epilepsia* 2008;49:425–430.
6. Menon R, Baheti NN, Cherian A, Iyer RS. Oxcarbazepine induced worsening of seizures in Jeavons syndrome: lessons learnt from an interesting presentation. *Neurol India* 2011; 59:70–72.

The Mystery Case series was initiated by the *Neurology*[®] Resident & Fellow Section to develop the clinical reasoning skills of trainees. Residency programs, medical student preceptors, and individuals were invited to use this Mystery Case as an educational tool. Responses were solicited through a group e-mail sent to the American Academy of Neurology Consortium of Neurology Residents and Fellows and through social media. All the answers that we received came through social media, from individuals rather than groups.

Most of the respondents (66%) correctly indicated Jeavons syndrome as the most likely diagnosis. The other preferred response was absence epilepsy. The most complete answer came from Dr. Felipe Borlot (Clinical Fellow, Toronto Western Hospital and University of Toronto, Canada). In his response, he pointed out that the key element for this Mystery Case is the fact that these reflex seizures are induced by eye closure and that the epileptiform abnormalities disappear with eye opening. The eyelid myoclonia in this patient is characteristic. Jeavons syndrome can be misdiagnosed as childhood or juvenile absence epilepsy, other forms of genetic or idiopathic generalized epilepsies, or even facial tics.

This Mystery Case illustrates a classic epilepsy syndrome, usually refractory to treatment, which persists throughout life.

Dragos A. Nita, MD, PhD, FRCPC
*Division of Neurology, The Hospital for Sick Children,
 University of Toronto, Canada*

Neurology®

Mystery Case: Eyelid myoclonia with absences in an adult patient

Yousef Hannawi, Shirish S. Satpute and Atul Maheshwari

Neurology 2014;82:e63-e64

DOI 10.1212/WNL.0000000000000139

This information is current as of February 24, 2014

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/82/8/e63.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2014/02/23/82.8.e63.DC1
References	This article cites 6 articles, 0 of which you can access for free at: http://n.neurology.org/content/82/8/e63.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Absence seizures http://n.neurology.org/cgi/collection/absence_seizures All clinical neurophysiology http://n.neurology.org/cgi/collection/all_clinical_neurophysiology All Epilepsy/Seizures http://n.neurology.org/cgi/collection/all_epilepsy_seizures EEG http://n.neurology.org/cgi/collection/eeg_ Epilepsy monitoring http://n.neurology.org/cgi/collection/epilepsy_monitoring_
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.