deficits may be due to an intact blood–brain barrier (BBB), which restricts entry of serum IgG into the CNS. It has recently been reported that AQP4-IgG can be present in serum several years before clinical onset of NMO.⁷ In addition, experimental studies have shown induction of NMO-like histopathology by transfer of human AQP4-IgG to animals only provided BBB breakdown.²

Further studies are needed to evaluate the neurologic outcome of infants and children of mothers with NMO given the evidence that AQP4-IgG can be passively transferred to the fetus.

From Vejle Hospital (N.A.); Institute of Molecular Medicine (N.A.), University of Southern Denmark, Odense; Aarhus University Hospital Skejby (T.B.H.); Aarhus University Hospital (T.P.); Odense University Hospital (S.T.L.), Denmark; and Mayo Clinic (B.G.W.), Rochester, MN.

Author contributions: Nasrin Asgari: study concept and design, acquisition of data, interpretation of results, and writing of manuscript. T.B. Henriksen: clinical investigations, revising manuscript, and approving final version. T. Petersen: acquisition of data, revising manuscript, and approving final version. S.T. Lillevang: determination of aquaporin-4 antibodies, revising manuscript, and approving final version. B.G. Weinschenker: study concept and design, interpretation of result, revising manuscript, and approving final version, study supervisor.

Study funding: Supported by the Lundbeck Foundation, The Region of Southern Denmark, and The University of Southern Denmark.

Disclosure: N. Asgari has received a research grant from Lundbeck Foundation. T. Henriksen, T. Petersen, and S. Lillevang report no disclosures relevant to the manuscript. B. Weinschenker has received a research grant from the Guthy Jackson Foundation. He receives royalties from RSR for a technology license related to a test for aquaporin-4 autoantibodies for diagnosis of neuromyelitis optica. He serves on

data safety monitoring committees for Novartis, Biogen-Idec, and Mitsubishi pharmaceutical companies, and serves on an adjudication panel for Medimmune Pharmaceuticals. He served as a consultant for GlaxoSmithKline, Elan, Ono, Chugai, and Alexion pharmaceutical companies. He serves on editorial boards for Neurology®, Canadian Journal of Neurological Sciences, and Turkish Journal of Neurology. Go to Neurology.org for full disclosures.

Received February 18, 2014. Accepted in final form May 22, 2014. Correspondence to Dr. Asgari: nasgari@health.sdu.dk

© 2014 American Academy of Neurology

1. Wingerchuk DM, Lennon VA, Lucchinetti CF, Pittock SJ, Weinschenker BG. The spectrum of neuromyelitis optica. *Lancet Neurol* 2007;6:805–815.
2. Bradl M, Lassmann H. Experimental models of neuromyelitis optica. *Brain Pathol* 2014;24:74–82.
3. Saadoun S, Waters P, Leite MI, Bennett JL, Vincent A, Papadopoulos MC. Neuromyelitis optica IgG causes placental inflammation and fetal death. *J Immunol* 2013;191:2999–3005.
4. Reuss R, Rommer PS, Bruck W, et al. A woman with acute myelopathy in pregnancy: case outcome. *BMJ* 2009;339:b4026.
5. Lefvert AK, Osterman PO. Newborn infants to myasthenic mothers: a clinical study and an investigation of acetylcholine receptor antibodies in 17 children. *Neurology* 1983;33:133–138.
6. Ringelstein M, Harmel J, Distelmaier F, et al. Neuromyelitis optica and pregnancy during therapeutic B cell depletion: infant exposure to anti-AQP4 antibody and prevention of rebound relapses with low-dose rituximab postpartum. *Mult Scler* 2013;19:1544–1547.
7. Nishiyama S, Ito T, Misu T, et al. A case of NMO seropositive for aquaporin-4 antibody more than 10 years before onset. *Neurology* 2009;72:1960–1961.

Juan D. Ramirez, MD
Abdella M. Habib, PhD
James J. Cox, PhD
Andreas C.

Themistocleous,
MBBCh, PhD, FCP
Stephen B. McMahon
PhD, FMedSci, FSB
John N. Wood, PhD,
FRS
David L.H. Bennett
PhD, FRCP

NULL MUTATION IN SCN9A IN WHICH NOXIOUS STIMULI CAN BE DETECTED IN THE ABSENCE OF PAIN

OPEN

The term nociception was originally defined by Sherrington¹ as the neural process by which high-threshold stimuli (which cause tissue injury) are detected. This is distinct from pain, which is defined by the quality of the sensory percept (i.e., unpleasant), is not always evoked by a noxious stimulus, and can also be experienced in the absence of a noxious stimulus. In man, recessive loss of function mutations in the *SCN9A* gene, encoding the α subunit of $\text{Na}_v1.7$, results in anosmia and the congenital inability to experience pain,^{2–4} but there have been few reports of detailed sensory testing in these patients.

Case report. A 27-year-old man who participated in the Painful Channelopathies Study (NRES-UK reference: 12/LO/0017) had a history of self-mutilating behavior during childhood and multiple

painless injuries with normal cognitive development. He had never perceived pain but reported that otherwise somatosensation was normal. In his late teens, he learned to avoid injury by attending to what he described as a tingling sensation, which was not unpleasant but that he noted only occurred in the context of threatened tissue injury. The family pedigree is shown in the figure, A. He had 2 sisters, one of whom died in childhood as a consequence of sepsis. The surviving sister also has congenital inability to experience pain.

The proband presented with short stature of 149 cm. He was anosmic. Tone and power were normal throughout and all deep tendon reflexes were preserved. He had no impairment in light touch, vibration detection, or proprioception. Pinprick was not perceived as painful although he could detect that it was pointed and a flare developed shortly after stimulation.

Genetic testing for *SCN9A* mutations in the index case and his sister (figure, A) revealed compound heterozygous mutations (figure, B). Nerve conduction

Supplemental data
at Neurology.org

Figure Detailed sensory phenotyping in a patient with compound heterozygous null mutations in $Na_v1.7$

(A) The family pedigree shows the proband (arrow) and 2 sisters with congenital inability to experience pain. (B) Cartoon shows the structure of $Na_v1.7$ showing the typical 4-domain structure of voltage-gated sodium channels each with 6 transmembrane segments. The mutations are as follows: a premature stop codon at arginine 830 (within domain 2) and a frameshift at glutamate 1773 within the C-terminal domain. (C) Graphical representation of sensory testing in the proband expressed as Z-scores: cold detection threshold (CDT), warm detection threshold (WDT), thermal sensory limen (TSL), cold pain threshold (CPT), heat pain threshold (HPT), pressure pain threshold (PPT), mechanical pain threshold (MPT), mechanical pain sensitivity (MPS), wind-up ratio (WUR), mechanical detection threshold (MDT), and vibration detection threshold (VDT). (D) Graph represents the patient's rating of his tingling sensation on a numerical rating scale (means $\pm$ SE) in response to suprathreshold thermal stimuli delivered in a randomized manner. The threshold for this distinct sensation is clearly in the noxious range and the intensity of the sensation encodes the strength of the thermal stimulus. (E) Photomicrograph of skin section immunostained for PGP 9.5 (red) to demonstrate nerve fibers and collagen type IV (green) to show the basement membrane. Fibers are clearly seen in the dermal plexus and some are crossing into the epidermis (arrows); however, the intraepidermal nerve fiber density is reduced to 3.98 fibers/mm. Scale bar: 50 μ m.

studies demonstrated small-amplitude sural sensory nerve action potentials (table e-1 on the *Neurology*[®] Web site at Neurology.org). Quantitative sensory testing was performed according to the German Neuropathic Pain Network⁵ (e-Methods). He did not experience pain (i.e., there was no unpleasant sensation) in response to any stimulus applied including extremes of temperature or mechanical stimuli. His mechanical detection and vibration detection thresholds were normal. He was hyposensitive to warm and cool stimuli (figure, C). High temperatures and strong mechanical stimuli evoked a mild tingling sensation. This sensation was never unpleasant but he had learned to use it as an injury signal. We therefore performed suprathreshold thermal stimulation by giving randomized thermal stimuli and asking him to rate the intensity of this sensation (figure, D). The threshold for this stimulus was in the noxious range at 42°C and the intensity of the sensation could encode the strength of the stimulus. Skin biopsy taken from the distal leg showed that intraepidermal nerve fibers were present although at a density below the lower limit of normal (figure, E).

Discussion. The index case reported in this kindred demonstrated insensitivity to pain, normal large-fiber sensory function, and anosmia: typical features of congenital insensitivity to pain (CIP) secondary to *SCN9A* mutations. Genetic testing revealed compound heterozygous mutations in *SCN9A*: a premature stop codon in coding exon 15 (c.2488C>T), which has previously been reported,³ and a previously unreported 1-bp deletion within coding exon 26 (c.5318delA), which induces a frameshift at position 1773 in the C-terminal domain of the channel.

The patient had identified a sensation that was not unpleasant but only occurred in the context of strong mechanical or thermal stimuli that could cause tissue injury. The sensation was mild, localized to the stimulus, and had a tingling quality to it. He used this sensation as a warning signal and had managed to reduce the frequency of injury. It is currently unclear as to the mechanism by which he can detect tissue injury in the absence of pain. Nociceptors still innervate the skin, albeit at a lower density than in age-matched controls. The proposed mechanisms through which null mutations in *Nav1.7* result in inability to experience pain include impaired transduction in nociceptor terminals, action potential transmission along the axon, and release of neurotransmitter from central terminals of sensory afferents.^{6,7} It is possible that a small population of nociceptive afferents are able to function in the absence of *Nav1.7* but provide insufficient input to drive pain sensation. This case further emphasizes that the detection of noxious stimuli and the sensation of pain are distinct and could also

have practical implications if other patients with CIP can train themselves to attend to sensations that may indicate potential tissue injury but are not painful.

From the University of Oxford (J.D.R., A.C.T., D.L.H.B.); Wolfson Institute for Biomedical Research (A.M.H., J.J.C., J.N.W.), University College London; and Wolfson CARD (S.B.M.), King's College London, UK.

Author contributions: Dr. D.L. Bennett: study concept and design, analysis and interpretation, study supervision, critical revision of the manuscript for important intellectual content. Dr. S.B. McMahon: study concept and design, analysis and interpretation, critical revision of the manuscript for important intellectual content. Dr. J.N. Wood: analysis and interpretation, critical revision of the manuscript for important intellectual content. J.D. Ramirez: study concept and design, acquisition of data, analysis and interpretation, generating first draft of manuscript. A.M. Habib: acquisition of data, analysis and interpretation. A. Themistocleous: acquisition of data, analysis and interpretation. J.J. Cox: acquisition of data, analysis and interpretation, critical revision of the manuscript for important intellectual content.

Study funding: Supported by a strategic award by the Wellcome Trust to the London Pain Consortium (ref. no. 083259) and a Medical Research Council career development fellowship.

Disclosure: J. Ramirez is sponsored by a Francisco Jose de Caldas Scholarship from Colciencias, through LASPAU, Harvard University. He was also funded by the Wellcome Trust-funded London Pain Consortium (ref. no. 083259). A. Habib is funded by an MRC Research Career Development fellowship. J. Cox is funded by an MRC Research Career Development fellowship. A. Themistocleous is funded by the Wellcome Trust-funded London Pain Consortium (ref. no. 083259). S. McMahon is a member of the Innovative Medicines Initiative Europain (grant 115007) and has acted as a consultant for Pfizer, Astra Zeneca, Mundipharma, and Bayer. He has received grant funding support from Astra Zeneca, Pfizer, and Covergence. He is a Wellcome Trust senior investigator and a member of the Wellcome Trust-funded London Pain Consortium (ref. no. 083259). J. Wood is a member of the Innovative Medicines Initiative Europain (grant 115007). He is also funded by the Medical Research Council, the BBSRC, is a Wellcome Trust senior investigator and a member of the Wellcome Trust-funded London Pain Consortium (ref. no. 083259). D. Bennett is a member of the Innovative Medicines Initiatives (public-private partnerships between the EU and EFPLA), Europain (grant 115007), and Stembancc. He has acted as a consultant for Pfizer. David Bennett is a senior Wellcome Trust Clinical Scientist Fellow (ref. no. 095698z/11/z) and a member of the Wellcome Trust-funded London Pain Consortium (ref. no. 083259). Go to Neurology.org for full disclosures. The Article Processing Charge was paid by the Wellcome Trust.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received March 28, 2014. Accepted in final form June 12, 2014.

Correspondence to Dr. Bennett: david.bennett@ndcn.ox.ac.uk

© 2014 American Academy of Neurology

1. Sherrington CS. Qualitative differences of spinal reflex corresponding with qualitative difference of cutaneous stimulus. *J Physiol* 1903;30:39–46.
2. Cox JJ, Reimann F, Nicholas AK, et al. An *SCN9A* channelopathy causes congenital inability to experience pain. *Nature* 2006;444:894–898.
3. Goldberg YP, MacFarlane J, MacDonald ML, et al. Loss-of-function mutations in the *Nav1.7* gene underlie congenital indifference to pain in multiple human populations. *Clin Genet* 2007;71:311–319.

4. Ahmad S, Dahllund L, Eriksson AB, et al. A stop codon mutation in SCN9A causes lack of pain sensation. *Hum Mol Genet* 2007;16:2114–2121.
5. Rolke R, Baron R, Maier C, et al. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. *Pain* 2006;123:231–243.
6. Raouf R, Rugiero F, Kiesewetter H, et al. Sodium channels and mammalian sensory mechanotransduction. *Mol Pain* 2012;8:21.
7. Minett MS, Nassar MA, Clark AK, et al. Distinct Nav1.7-dependent pain sensations require different sets of sensory and sympathetic neurons. *Nat Commun* 2012; 3:791.

Subspecialty Alerts by E-mail!

Customize your online journal experience by signing up for e-mail alerts related to your subspecialty or area of interest. Access this free service by visiting Neurology.org/site/subscriptions/etoc.xhtml or click on the “E-mail Alerts” link on the home page. An extensive list of subspecialties, methods, and study design choices will be available for you to choose from—allowing you priority alerts to cutting-edge research in your field!

Your Research Recognized: Apply for an AAN Award

Have you, or someone you know, been conducting research that may be advancing the field of neurology? Apply—or nominate a colleague—for a prestigious AAN award and your research could be recognized during the 2015 AAN Annual Meeting in Washington, DC, April 18–25, 2015, the world’s largest gathering of neurologists. Awards are available for all career stages and interests. View all 2015 AAN awards and application deadlines at AAN.com/view/15Awards. Application deadline is October 23, 2014.

WriteClick® rapid online correspondence

The editors encourage comments about recent articles through WriteClick:

Go to *Neurology.org* and click on the “WriteClick” tab at the top of the page. Responses will be posted within 72 hours of submission.

Before using WriteClick, remember the following:

- WriteClick is restricted to comments about studies published in *Neurology* within the last eight weeks
- Read previously posted comments; redundant comments will not be posted
- Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
- You can include a maximum of five authors (including yourself)

Neurology®

Null mutation in *SCN9A* in which noxious stimuli can be detected in the absence of pain

Juan D. Ramirez, Abdella M. Habib, James J. Cox, et al.

Neurology 2014;83;1577-1580 Published Online before print September 24, 2014

DOI 10.1212/WNL.0000000000000913

This information is current as of September 24, 2014

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/83/17/1577.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2014/09/24/WNL.0000000000000913.DC1
References	This article cites 7 articles, 0 of which you can access for free at: http://n.neurology.org/content/83/17/1577.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Pain http://n.neurology.org/cgi/collection/all_pain Ion channel gene defects http://n.neurology.org/cgi/collection/ion_channel_gene_defects Peripheral neuropathy http://n.neurology.org/cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.