Teaching *NeuroImages*: Reversible pontomesencephalic edema caused by traumatic carotid cavernous fistula

Yazan J. Alderazi,
MBBCh
Sushrut Dharmadhikari,
MD
Diogo C. Haussen, MD
Dileep R. Yavagal, MD

Correspondence to
Dr. Yavagal:
DYavagal@med.miami.edu

Figure Pretreatment pontomesencephalic lesion

(A) Fluid-attenuated inversion recovery and (B) T1 postcontrast MRI. (C) Carotid-cavernous fistula drainage via bridging veins (arrow) to the anterior pontomesencephalic vein (interrupted arrow) and via superior ophthalmic vein (arrowhead) and pterygoid plexus (double arrows) on catheter angiography, lateral view. (D) Pontomesencephalic edema resolution on post-treatment MRI.

A 45-year-old man presented with a painful bulging right eye and blindness 6 weeks after a motorcycle accident and traumatic brain injury. He had a complete right oculomotor nerve palsy and only perception of light. MRI revealed ipsilateral pontomesencephalic edema (figure, A and B). Catheter angiography showed a direct right carotid-cavernous fistula (figure, C). Two months after coil embolization, the ptosis, proptosis and ophthalmoplegia resolved. MRI revealed resolution of pontomesencephalic edema (figure, D). The presumed mechanism was transmission of arterialized pressures of the cavernous sinus via a bridging vein to the anterior pontomesencephalic vein, generating venous hypertension and vasogenic edema.^{1,2}

AUTHOR CONTRIBUTIONS

Study concept and design: Y.J.A., D.Y. Acquisition of data: Y.J.A. Analysis and interpretation of data: Y.J.A., S.D., D.C.H. Drafting of the

manuscript: Y.J.A., S.D. Critical revision of the manuscript for important intellectual content: S.D., D.C.H., D.Y. Study supervision: D.Y.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

REFERENCES

1. Kiyosue H, Tanoue S, Sagara Y, et al. The anterior medullary-anterior pontomesencephalic venous system and its bridging veins communicating to the dural sinuses: normal anatomy and drainage routes from dural arteriovenous fistulas. *Neuroradiology* 2008;50:1013–1023.
2. Kiyosue H, Mori H, Sagara Y, et al. Basal cerebral venous drainage from cavernous sinus dural arteriovenous fistulas. *Neuroradiology* 2009;51:175–181.

Neurology®

Teaching *NeuroImages*: Reversible pontomesencephalic edema caused by traumatic carotid cavernous fistula

Yazan J. Alderazi, Sushrut Dharmadhikari, Diogo C. Haussen, et al.

Neurology 2014;83:e18

DOI 10.1212/WNL.0000000000000562

This information is current as of July 7, 2014

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/83/2/e18.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2014/07/06/83.2.e18.DC1
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/83/2/e18.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Carotid artery dissection http://n.neurology.org/cgi/collection/carotid_artery_dissection
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2014 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.